



AP50 / AP2M1 Rabbit anti-Human Polyclonal (aa1-150) Antibody - LS-C185408 - LSBio	
CatalogID:	LS-C185408
Target:	adaptor-related protein complex 2, mu 1 subunit (AP2M1)
Synonyms:	AP2M1 Antibody, Adaptin-mu2 Antibody, AP-2 complex subunit mu Antibody, AP-2 mu 2 chain Antibody, AP-2 mu chain Antibody, AP50 Antibody, CLAPM1 Antibody, Mu2 Antibody, KIAA0109 Antibody, HA2 50 kDa subunit Antibody
Host	AP2M1 antibody was produced in Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Immunogen Species:	AP50 / AP2M1 antibody was raised against Human
Antigen Type:	Recombinant protein
Immunogen:	AP50 / AP2M1 antibody was raised against recombinant protein fragment contain a sequence corresponding to a region within amino acids 1 and 150 (Q96CW1) of AP2M1. Percent identity by BLAST analysis: Human, Mouse, Rat, Bovine (100%); Pig, Xenopus (99%); Chicken (97%).
Specificity:	Human AP50 / AP2M1
Epitope:	aa1-150
Reactivity:	Human, Mouse, Rat, Bovine
Predicted Reactivity:	Pig, Chicken, Xenopus
Purification:	Immunoaffinity purified
Presentation:	0.1 M Tris-glycine, pH 7.0, 10% glycerol, 0.01% Thimerosal
Recommended Storage:	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Usage Summary:	IHC-paraffin: Suggested antigen retrieval using heat mediated 10 mM Citrate buffer (pH 6.0) or Tris-EDTA buffer (pH 8.0).
Uses:	IHC - Paraffin (1:100 - 1:1000), Western blot (1:500 - 1:3000) (Optimal dilution to be determined by the researcher)
Size:	100 µl
Concentration:	0.57 mg/ml

Immunohistochemistry Image:



IHC of paraffin-embedded N87 xenograft using AP2M1 antibody at 1:100 dilution.

Western Blot Image:



Sample (30 ug of whole cell lysate). A:293T, B: HeLa S3. 10% SDS PAGE. AP50 / AP2M1 antibody diluted at 1:500 $\,$

Requested From: Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/26/2014
© 2014 LifeSpan BioSciences