

CatalogID:	LS-C204411
Target:	CD82 molecule
Synonyms:	CD82 Antibody, 4F9 Antibody, C33 Antibody, C33 antigen Antibody, CD82 molecule Antibody, GR15 Antibody, IA4 Antibody, Inducible membrane protein R2 Antibody, KAI1 Antibody, Kangai 1 Antibody, R2 Antibody, R2 leukocyte antigen Antibody, SAR2 Antibody, ST6 Antibody, Tetraspanin-27 Antibody, Tspan-27 Antibody, TSPAN27 Antibody, CD82 antigen Antibody, Leukocyte surface antigen r2 Antibody, Metastasis suppressor Kangai-1 Antibody
Family / Subfamily:	Tetraspan / not assigned-Tetraspan
Host	CD82 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG2a
Clone Name:	C33
Conjugations:	Fluorescein (FITC)
Immunogen Species:	CD82 antibody was raised against Human
Antigen Type:	Cells
Immunogen:	CD82 antibody was raised against c91/PL (human HTLV-1+ T cell line).
Specificity:	Monoclonal antibody C33 reacts with CD82, a widely expressed cell surface protein of the tetraspanin family. CD82 is also found in endosome/lysosome compartments.
Reactivity:	Human
Purification:	Protein A purified
Presentation:	PBS, 0.2% BSA, 0.02% sodium azide.
Recommended Storage:	Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.
Usage Summary:	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan
La	boratory Reagent For In Vitro Research Use Only
Not for resale	without prior written consent from LifeSpan BioSciences, Inc.
	Created on 9/27/2014 © 2014 LifeSpan BioSciences