

45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope: aa1-22  Reactivity: Human  Purification: Protein A purified  Presentation: PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage: Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.		
Target: chemokine (C-C motif) receptor 5 (gene/pseudogene) (CCR5)  Synonyms: CCR5 Antibody, C-C chemokine receptor 5 Antibody, CD195 Antibody, CD195 Antibody, C-C CKR-5 Antibody, C-C chemokine receptor 5 Antibody, Chemokine c-c motif receptor 5 Antibody, Chemokine c-c motif receptor 5 Antibody, CMCKR5 Antibody, CKR5 Antibody, CCR5 Antibody, Chemokine receptor 5 Antibody, CCKR5 Antibody, CCR5 Antibody, CCR6 Antibody, CCR7 Antibody, CCR6 Antibody, CCR7 Antibody was produced in Mouse  Clonality: Monoclonal  Isotype: IgG1  Clone Name: T21/8  Conjugations: Phycoerythrin (PE)  Immunogen Species: CCR5 antibody was raised against Human  Antigen Type: Synthetic peptide  Immunogen: CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity: Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 KD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope: aa1-22  Reactivity: Human  Purification: Protein A purified  Presentation: Pss, 0.2% BSA, 0.02% sodium azide.  Recommended Storage: Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.	CCR5 Mouse anti-Humar	Monoclonal (aa1-22) (PE) (T21/8) Antibody - LS-C204488 - LSBio
Synonyms:  CCR5 Antibody, C-C chemokine receptor 5 Antibody, CD195 antigen Antibody, CD195 Antibody, C-C CKR-5 Antibody, Chemokine c-c motif receptor 5 Antibody, Chemria Antibody, CMCKR5 Antibody, CC chemokine receptor 5 Antibody, CMCKR5 Antibody, CCF chemokine receptor 5 Antibody, CMCKR5 Antibody, CCF chemokine receptor Antibody, CHEMPA Antibody, CCCKR5 Antibody, CCF chemokine receptor Antibody, CCR-5 Antibody, CCC-CKR-5 Antibody, CKR-5 Antibody, CCC-CKR-5 Antibody, CKR-5 Antibody, CCC-CKR-5 Antibody, CKR-5 Antibody, CKR-5 Antibody, CCC-CKR-5 Antibody, CKR-5 Antibody, CCC-CKR-5 Antibody, CKR-5 Antibody, CCC-CKR-5 Antibody, CKR-5 Antibody, CCC-CKR-5 Antibody, CKR-5 Antibody was produced in Mouse  Clonality:  Monoclonal  Isotype:  IgG1  Clone Name:  C21/8  Conjugations:  Phycoerythrin (PE)  Immunogen Species:  CCR5 antibody was raised against Human  Antigen Type:  Synthetic peptide  Immunogen:  CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity:  Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 KD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope:  aa1-22  Reactivity:  Human  Purification:  Protein A purified  Presentation:  Protein A purified  Presentation:  Protein A purified  Presentation:  Presentation:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.	CatalogID:	LS-C204488
CD195 Antibody, C-C CKR-5 Antibody, Chemokine c-c mutif receptor 5 Antibody, CKRS Antibody, CKRS Antibody, CCRS Antibody, CCR-5 Antibody was produced in Mouse  Family / Subfamily:  GPCR / Chemokine  CCR5 antibody was produced in Mouse  Clonality:  Monoclonal  Isotype:  IgG1  Clone Name:  T21/8  Conjugations:  Phycoerythrin (PE)  Immunogen Species:  CCR5 antibody was raised against Human  Antigen Type:  Synthetic peptide  Immunogen:  CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity:  Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope:  aa1-22  Reactivity:  Human  Purification:  Protein A purified  Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 mil) is sufficient for 100 tests.  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Target:	chemokine (C-C motif) receptor 5 (gene/pseudogene) (CCR5)
Host Clonality: Monoclonal  Isotype: IgG1 Clone Name: T21/8 Conjugations: Phycoerythrin (PE) Immunogen Species: CCR5 antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity: Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope: aa1-22 Reactivity: Human Purification: Protein A purified Presentation: PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage: Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary: The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses: Flow Cytometry (Optimal dilution to be determined by the researcher) Size: 100 tst	Synonyms:	CD195 Antibody, C-C CKR-5 Antibody, Chemokine c-c motif receptor 5 Antibody, Chemr13 Antibody, CMKBR5 Antibody, CKR5 Antibody, Cc chemokine receptor 5 Antibody, CCCKR5 Antibody, CCR-5 Antibody, Ccr5 receptor Antibody, Chemokine receptor CCR5 Antibody, HIV-1 fusion coreceptor Antibody, IDDM22 Antibody, C-C chemokine receptor type 5 Antibody, CC-CKR-5 Antibody, CKR-5
Clonality: Monoclonal Isotype: IgG1 Clone Name: T21/8 Conjugations: Phycoerythrin (PE) Immunogen Species: CCR5 antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate. Specificity: Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 KD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope: aa1-22 Reactivity: Human Purification: Protein A purified Presentation: PBS, 0.2% BSA, 0.02% sodium azide. Recommended Storage: Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light. Usage Summary: The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 mil) is sufficient for 100 tests.  Uses: Flow Cytometry (Optimal dilution to be determined by the researcher) Size: 100 tst	Family / Subfamily:	GPCR / Chemokine
Isotype: IgG1  Clone Name: T21/8  Conjugations: Phycoerythrin (PE)  Immunogen Species: CCR5 antibody was raised against Human  Antigen Type: Synthetic peptide  Immunogen: CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity: Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope: aa1-22  Reactivity: Human  Purification: Protein A purified  Presentation: PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage: Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary: The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 mil) is sufficient for 100 tests.  Uses: Flow Cytometry (Optimal dilution to be determined by the researcher)  Size: 100 tst	Host	CCR5 antibody was produced in Mouse
Clone Name:  Conjugations: Phycoerythrin (PE)  Immunogen Species: CCR5 antibody was raised against Human  Antigen Type: Synthetic peptide  CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity: Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope: aa1-22  Reactivity: Human  Purification: Protein A purified  Presentation: PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage: Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary: The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses: Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Clonality:	Monoclonal
Conjugations:    Phycoerythrin (PE)	Isotype:	IgG1
Immunogen Species:  CCR5 antibody was raised against Human  Synthetic peptide  CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity:  Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope:  aa1-22  Reactivity:  Human  Purification:  Protein A purified  Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Clone Name:	T21/8
Antigen Type:  Immunogen:  CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.  Specificity:  Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope:  aa1-22  Reactivity:  Human  Purification:  Protein A purified  Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Conjugations:	Phycoerythrin (PE)
Immunogen:       CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.         Specificity:       Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.         Epitope:       aa1-22         Reactivity:       Human         Protein A purified       Presentation:         PBS, 0.2% BSA, 0.02% sodium azide.         Recommended Storage:       Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.         Usage Summary:       The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.         Uses:       Flow Cytometry (Optimal dilution to be determined by the researcher)         Size:       100 tst	Immunogen Species:	CCR5 antibody was raised against Human
Specificity:  Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope:  aa1-22  Reactivity:  Human  Protein A purified  Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Antigen Type:	Synthetic peptide
45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.  Epitope:  aa1-22  Reactivity:  Human  Protein A purified  Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Immunogen:	CCR5 antibody was raised against cCR5 peptide (Met1-Lys22) KLH conjugate.
Reactivity:  Human  Purification:  Protein A purified  Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Specificity:	Monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kD G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.
Purification:  Protein A purified  Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Epitope:	aa1-22
Presentation:  PBS, 0.2% BSA, 0.02% sodium azide.  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Reactivity:	Human
Recommended Storage:  Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Purification:	Protein A purified
Usage Summary:  The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses:  Flow Cytometry (Optimal dilution to be determined by the researcher)  Size:  100 tst	Presentation:	PBS, 0.2% BSA, 0.02% sodium azide.
ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.  Uses: Flow Cytometry (Optimal dilution to be determined by the researcher)  Size: 100 tst	Recommended Storage:	Store in the dark at 2-8 C. Do not freeze. Avoid prolonged exposure to light.
Size: 100 tst	Usage Summary:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 ul reagent / 100 ul of whole blood or 1 million cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
	Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Requested From: Japan	Size:	100 tst
	Requested From:	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/27/2014

© 2014 LifeSpan BioSciences