

CatalogID:	LS-C209604
Target:	protein C receptor, endothelial (PROCR)
Synonyms:	PROCR Antibody, Activated protein C receptor Antibody, CD201 Antibody, CD201 antigen Antibody, Centrocyclin Antibody, CCD41 Antibody, Endothelial protein C receptor Antibody, APC receptor Antibody, BA42O4.2 Antibody, CCCA Antibody, EPCR Antibody
Host	PROCR antibody was produced in Rat
Clonality:	Monoclonal
sotype:	lgG1
Clone Name:	RCR-252
Conjugations:	Fluorescein (FITC)
mmunogen Species:	PROCR / EPCR antibody was raised against Human
Antigen Type:	Cells
mmunogen:	PROCR / EPCR antibody was raised against human EPCR-positive RE-1 cells.
Specificity:	Human PROCR / EPCR
Reactivity:	Human
Purification:	Purified
Presentation:	PBS, 1% bovine serum
Recommended Storage:	Store at 4°C, stable for one year.
Usage Summary:	For flow cytometry and Western blot dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. For functional studies, in vitro dilutions have to be optimized by the user.
Uses:	Western blot, Immunoprecipitation, Flow Cytometry, Functional Assay (Optimal dilution to be determined by the researcher)
Size:	100 µg
Requested From:	Japan
La	aboratory Reagent For In Vitro Research Use Only
Not for resale	without prior written consent from LifeSpan BioSciences, Inc. Created on 9/27/2014