


LCN2 / Lipocalin 2 / NGAL Rabbit anti-Human Polyclonal (FITC) Antibody - LS-C211182 - LSBio

| | |
|-----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CatalogID: | LS-C211182 |
| Target: | lipocalin 2 (LCN2) |
| Synonyms: | LCN2 Antibody, 24p3 Antibody, HNL Antibody, Lipocalin 2 Antibody, Lipocalin-2 Antibody, MSF1 Antibody, Siderocalin LCN2 Antibody, NGAL Antibody, p25 Antibody, Lipocalin 2 (oncogene 24p3) Antibody, Oncogene 24p3 Antibody |
| Host | LCN2 antibody was produced in Rabbit |
| Clonality: | Polyclonal |
| Isotype: | IgG |
| Conjugations: | Fluorescein (FITC) |
| Immunogen Species: | LCN2 / Lipocalin 2 / NGAL antibody was raised against Human |
| Reactivity: | Human |
| Purification: | Caprylic acid and ammonium sulfate precipitation |
| Presentation: | PBS, pH 7.4, 0.03% Proclin 300, 50% glycerol. |
| Recommended Storage: | Aliquot and store at -20°C or -80°C. Avoid freeze-thaw cycles. |
| Uses: | IHC, Western blot, ELISA (Optimal dilution to be determined by the researcher) |
| Size: | 100 µg |

Western Blot Image:



Western blot of NGAL Antibody at 2 ug/ml + positive serum. Lane 1: positive serum at 1:10. Lane 2: positive serum at 1:50. Secondary: Goat polyclonal to Rabbit IgG at 1:15000 dilution. Predicted band size: 22 kDa. Observed band size: 22 kDa.

| | |
|------------------------------------------------------------------------------|-------|
| Requested From: | Japan |
| Laboratory Reagent For In Vitro Research Use Only | |
| Not for resale without prior written consent from LifeSpan BioSciences, Inc. | |
| Created on 9/27/2014 | |
| © 2014 LifeSpan BioSciences | |