

CCL2 / MCP-1 Rabbit anti-Human Polyclonal (FITC) Antibody - LS-C211505 - LSBio

| | |
|-----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CatalogID: | LS-C211505 |
| Target: | chemokine (C-C motif) ligand 2 (CCL2) |
| Synonyms: | CCL2 Antibody, C-C motif chemokine 2 Antibody, Chemokine (C-C motif) ligand 2 Antibody, GDCF-2 Antibody, HC11 Antibody, HSMCR30 Antibody, MCP1 Antibody, SCYA2 Antibody, Small-inducible cytokine A2 Antibody, Monocyte secretory protein JE Antibody, MCAF Antibody, MCP-1 Antibody, Monocyte chemotactic protein 1 Antibody, SMC-CF Antibody |
| Family / Subfamily: | Intercrine / Intercrine beta |
| Host | CCL2 antibody was produced in Rabbit |
| Clonality: | Polyclonal |
| Isotype: | IgG |
| Conjugations: | Fluorescein (FITC) |
| Immunogen Species: | CCL2 / MCP-1 antibody was raised against Human |
| Antigen Type: | Recombinant protein |
| Immunogen: | CCL2 / MCP-1 antibody was raised against recombinant human C-C motif chemokine 2 protein. |
| Specificity: | Human CCL2 / MCP-1 |
| Reactivity: | Human |
| Purification: | Caprylic acid and ammonium sulfate precipitation |
| Presentation: | PBS, pH 7.4, 0.03% Proclin 300, 50% glycerol. |
| Recommended Storage: | Aliquot and store at -20°C or -80°C. Avoid freeze-thaw cycles. |
| Uses: | IHC, Western blot, ELISA (Optimal dilution to be determined by the researcher) |
| Size: | 100 µg |
| Requested From: | Japan |

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/27/2014

© 2014 LifeSpan BioSciences