

**CST3 / Cystatin C Goat anti-Human Polyclonal (FITC) Antibody - LS-C212349 - LSBio**

|                             |  |
|-----------------------------|--|
| <b>CatalogID:</b>           | LS-C212349   |
| <b>Target:</b>              | cystatin C (CST3)  |
| <b>Synonyms:</b>            | CST3 Antibody, ARMD11 Antibody, BA218C14.4 (cystatin C) Antibody, Cystatin-3 Antibody, Cystatin 3 Antibody, Cystatin C Antibody, Cystatin-C Antibody, Gamma-trace Antibody, Post-gamma-globulin Antibody |
| <b>Host</b>                 | CST3 antibody was produced in Goat   |
| <b>Clonality:</b>           | Polyclonal   |
| <b>Isotype:</b>             | IgG  |
| <b>Conjugations:</b>        | Fluorescein (FITC)   |
| <b>Immunogen Species:</b>   | CST3 / Cystatin C antibody was raised against Human  |
| <b>Antigen Type:</b>        | Recombinant protein  |
| <b>Immunogen:</b>           | CST3 / Cystatin C antibody was raised against recombinant human Cystatin-C protein.  |
| <b>Specificity:</b>         | Human CST3 / Cystatin C  |
| <b>Reactivity:</b>          | Human  |
| <b>Purification:</b>        | Caprylic acid and ammonium sulfate precipitation   |
| <b>Presentation:</b>        | PBS.   |
| <b>Recommended Storage:</b> | Aliquot and store at -20°C or -80°C. Avoid freeze-thaw cycles.   |
| <b>Uses:</b>                | IHC, Western blot, ELISA (Optimal dilution to be determined by the researcher)   |
| <b>Size:</b>                | 100 µg   |
| <b>Requested From:</b>      | Japan  |

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/27/2014

© 2014 LifeSpan BioSciences