

| DDC / DOPA Decarboxylase Rabbit anti-Human Polyclonal (aa39-69) (FITC) Antibody - LS- C259175 - LSBio | |
|--|---|
| CatalogID: | LS-C259175 |
| Target: | dopa decarboxylase (aromatic L-amino acid decarboxylase) (DDC) |
| Synonyms: | DDC Antibody, AADC Antibody, DOPA decarboxylase Antibody |
| Family / Subfamily: | Carboxy-lyase / Not assigned-Other |
| Host | DDC antibody was produced in Rabbit |
| Clonality: | Polyclonal |
| Isotype: | IgG |
| Conjugations: | Fluorescein (FITC) |
| Immunogen Species: | DDC / DOPA Decarboxylase antibody was raised against Human |
| Antigen Type: | Synthetic peptide |
| Immunogen: | DDC / DOPA Decarboxylase antibody was raised against kLH-conjugated synthetic peptide (aa39-69) from the N-terminal region of human DDC. |
| Specificity: | Human DDC / DOPA Decarboxylase |
| Epitope: | aa39-69 |
| Reactivity: | Human, Mouse |
| Purification: | Affinity purified |
| Presentation: | PBS, pH 7.2, 0.09% sodium azide. |
| Recommended Storage: | May be stored at 4°C for short-term only. Aliquot to avoid repeated freezing and thawing. Store at -20°C. Aliquots are stable for 6 months. |
| Usage Summary: | The applications listed above are for the unconjugated form of this antibody. The conjugated antibody has not been tested. |
| Uses: | IHC, ICC, Western blot, Flow Cytometry, ELISA (Optimal dilution to be determined by the researcher) |
| Size: | 200 μΙ |
| Requested From: | Japan |

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 9/29/2014
© 2014 LifeSpan BioSciences