

Clonality: Polyclonal Isotype: IgG Conjugations: Fluorescein (FITC) Immunogen Species: TMIE antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size:	
Host TMIE antibody was produced in Rabbit Clonality: Polyclonal Isotype: IgG Conjugations: Fluorescein (FITC) Immunogen Species: TMIE antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Clonality: Polyclonal Isotype: IgG Conjugations: Fluorescein (FITC) Immunogen Species: TMIE antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Antibody
Isotype: IgG Conjugations: Fluorescein (FITC) Immunogen Species: TMIE antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Conjugations: Fluorescein (FITC) Immunogen Species: TMIE antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Reacommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Immunogen Species: TMIE antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Antigen Type: Synthetic peptide Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Immunogen: TMIE antibody was raised against kLH-conjugated synthetic from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
from internal region of human TMIE. Specificity: Human TMIE Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: 200 µl Japan Laboratory Reagent For In Vitro Research Use Only	
Epitope: aa78-108 Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	peptide (aa78-108)
Reactivity: Human Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form or conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: Size: 200 μl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Purification: Affinity purified Presentation: PBS, pH 7.2, 0.09% sodium azide. Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: Size: 200 μl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Presentation:PBS, pH 7.2, 0.09% sodium azide.Recommended Storage:May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months.Usage Summary:The applications listed above are for the unconjugated form of conjugated antibody has not been tested.Uses:Western blot, ELISA (Optimal dilution to be determined by the Size:Size:200 µlRequested From:JapanLaboratory Reagent For In Vitro Research Use Only	
Recommended Storage: May be stored at 4°C for short-term only. Aliquot to avoid rep thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form or conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
thawing. Store at -20°C. Aliquots are stable for 6 months. Usage Summary: The applications listed above are for the unconjugated form of conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: 200 µl Zapan Laboratory Reagent For In Vitro Research Use Only	
conjugated antibody has not been tested. Uses: Western blot, ELISA (Optimal dilution to be determined by the Size: Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	eated freezing and
Size: 200 µl Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	of this antibody. The
Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	e researcher)
Laboratory Reagent For In Vitro Research Use Only	
· · · · · · · · · · · · · · · · · · ·	
Not for resale without prior written consent from LifeSpan RioScience	
Created on 9/30/2014	es, Inc.