

rous sclerosis 2 (TSC2) 2 Antibody, LAM Antibody, Tuberous sclerosis 2 protein Antibody, TSC4 body, Tuberin Antibody, Tuberous sclerosis 2 Antibody 2 antibody was produced in Rabbit clonal boerythrin (PE) erin / TSC2 antibody was raised against Rat hetic phospho-peptide erin / TSC2 antibody was raised against kLH-conjugated synthetic phopeptide around Thr1373 of rat TSC2.
andy, Tuberin Antibody, Tuberous sclerosis 2 Antibody 2 antibody was produced in Rabbit clonal coerythrin (PE) erin / TSC2 antibody was raised against Rat hetic phospho-peptide erin / TSC2 antibody was raised against kLH-conjugated synthetic
clonal coerythrin (PE) erin / TSC2 antibody was raised against Rat hetic phospho-peptide erin / TSC2 antibody was raised against kLH-conjugated synthetic
oerythrin (PE) erin / TSC2 antibody was raised against Rat hetic phospho-peptide erin / TSC2 antibody was raised against kLH-conjugated synthetic
erin / TSC2 antibody was raised against Rat hetic phospho-peptide erin / TSC2 antibody was raised against kLH-conjugated synthetic
erin / TSC2 antibody was raised against Rat hetic phospho-peptide erin / TSC2 antibody was raised against kLH-conjugated synthetic
hetic phospho-peptide erin / TSC2 antibody was raised against kLH-conjugated synthetic
rin / TSC2 antibody was raised against kLH-conjugated synthetic
rin / TSC2 antibody was raised against kLH-conjugated synthetic phopeptide around Thr1373 of rat TSC2.
Fuberin / TSC2
1373
Rabbit
ity purified
, pH 7.2, 0.09% sodium azide.
be stored at 4°C for short-term only. Aliquot to avoid repeated freezing and ing. Store at -20°C. Aliquots are stable for 6 months.
applications listed above are for the unconjugated form of this antibody. The ugated antibody has not been tested.
A, Dot Blot (Optimal dilution to be determined by the researcher)
l
n
Reagent For In Vitro Research Use Only