


NANOG Rabbit anti-Mouse Polyclonal (C-Terminus) Antibody - LS-C287289 - LSBio	
<b>CatalogID:</b>	LS-C287289
<b>Target:</b>	Nanog homeobox
<b>Synonyms:</b>	NANOG Antibody, Nanog homeobox Antibody, NANOGP8 Antibody, HNanog Antibody, Homeobox protein NANOG Antibody
<b>Host</b>	NANOG antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Immunogen Species:</b>	NANOG antibody was raised against Mouse
<b>Specificity:</b>	Region between residue 250 and the C-terminus (residue 305) of mouse Nanog homeobox using the numbering given in TrEMBL entry Q80Z64 (GeneID 71950).
<b>Epitope:</b>	C-Terminus
<b>Reactivity:</b>	Mouse
<b>Purification:</b>	Immunoaffinity purified
<b>Presentation:</b>	Tris-citrate/phosphate buffer, pH 7-8, 0.09% sodium azide.
<b>Recommended Storage:</b>	Store at 2-8°C for up to 1 year.
<b>Uses:</b>	Western blot (1:1000 - 1:10000), Immunoprecipitation (2 - 5 µg/mg lysate) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 µl
<b>Manufacturer:</b>	Bethyl Laboratories, Inc.

**Western Blot Image:**



Detection of Mouse Nanog by Western Blot and Immunoprecipitation. Samples: Whole cell lysate (5, 15 and 50 µg for WB; 1 mg for IP, 20% of IP loaded) from mouse F9 cells. Antibodies: Affinity purified rabbit anti-Nanog antibody used for WB at 0.1 µg/ml (A) and 1 µg/ml (B) and used for IP at 3 µg/mg lysate. Nanog was also immunoprecipitated by a previous lot of this antibody. Detection: Chemiluminescence with exposure times of 30 seconds (A) and 10 seconds (B).

**Requested From:**

Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 9/30/2014

© 2014 LifeSpan BioSciences