

CD71 / Transferrin Receptor Rat anti-Mouse Monoclonal (RPE) (YTA74.4) Antibody - LS-C43243 - LSBio	
<b>CatalogID:</b>	LS-C43243
<b>Target:</b>	transferrin receptor (TFRC)
<b>Synonyms:</b>	TFRC Antibody, CD71 Antibody, TFR1 Antibody, TRFR Antibody, TFR Antibody, TR Antibody, CD71 antigen Antibody, p90 Antibody, T9 Antibody, Transferrin receptor Antibody, Transferrin receptor protein 1 Antibody
<b>Family / Subfamily:</b>	Protease / Metallopeptidase M28B
<b>Host</b>	TFRC antibody was produced in Rat
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG2a
<b>Clone Name:</b>	YTA74.4
<b>Conjugations:</b>	R. Phycoerythrin (RPE)
<b>Immunogen Species:</b>	CD71 / Transferrin Receptor antibody was raised against Mouse
<b>Antigen Type:</b>	Cells
<b>Immunogen:</b>	CD71 / Transferrin Receptor antibody was raised against concanavalin A activated mouse spleen cells.
<b>Specificity:</b>	Recognizes the murine CD71 cell surface antigen, expressed by all dividing cells. CD71 acts as a receptor for transferrin
<b>Reactivity:</b>	Mouse
<b>Purification:</b>	Protein G purified
<b>Reconstitution:</b>	Distilled Water.
<b>Presentation:</b>	Lyophilized, PBS, pH 7.4, 0.09% sodium azide, 1% BSA.
<b>Usage Summary:</b>	Flow Cytometry: Use 10 ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100 ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors.
<b>Uses:</b>	Flow Cytometry (1:1) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 tst
<b>Requested From:</b>	Japan
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
Created on 10/1/2014	
© 2014 LifeSpan BioSciences	