

**ICAM1 / CD54 Mouse anti-Human Monoclonal (RPE) Antibody - LS-C43797 - LSBio**

<b>CatalogID:</b>	LS-C43797
<b>Target:</b>	intercellular adhesion molecule 1 (ICAM1)
<b>Synonyms:</b>	ICAM1 Antibody, BB2 Antibody, CD54 antigen Antibody, CD54 Antibody, ICAM-1 Antibody, p3.58 Antibody
<b>Family / Subfamily:</b>	ICAM / not assigned-ICAM
<b>Host</b>	ICAM1 antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Conjugations:</b>	R. Phycoerythrin (RPE)
<b>Immunogen Species:</b>	ICAM1 / CD54 antibody was raised against Human
<b>Antigen Type:</b>	Cells
<b>Immunogen:</b>	ICAM1 / CD54 antibody was raised against human monocytes.
<b>Specificity:</b>	Recognizes the human CD54 cell surface antigen, a 90kD glycoprotein also known as ICAM-1. CD54 is expressed by many cells following activation by inflammatory mediators. Clone 15.2 is reported to block CD54 function.
<b>Reactivity:</b>	Human, Pig
<b>Purification:</b>	Purified
<b>Reconstitution:</b>	Distilled Water.
<b>Presentation:</b>	Lyophilized, PBS, pH 7.4, 1% BSA, 0.09% sodium azide.
<b>Usage Summary:</b>	Flow Cytometry: Use 10 ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100 ul. Method sheets are available on request.
<b>Uses:</b>	Flow Cytometry (1:1) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 tst
<b>Requested From:</b>	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 10/1/2014

© 2014 LifeSpan BioSciences