

FCGR1A / CD64 Mouse anti-Human Monoclonal (RPE) (10.1) Antibody - LS-C45349 - LSBio	
<b>CatalogID:</b>	LS-C45349
<b>Target:</b>	Fc fragment of IgG, high affinity Ia, receptor (CD64) (FCGR1A)
<b>Synonyms:</b>	FCGR1A Antibody, CD64A Antibody, CD64 Antibody, CD64 antigen Antibody, FCG1 Antibody, FcgammaR1a Antibody, IGFR1 Antibody, IgG Fc receptor I Antibody, Fc-gamma receptor I A1 Antibody, Fc-gamma R1A Antibody, FCGR1 Antibody, Fc gamma receptor Antibody, Fc-gamma RI Antibody, FCRI Antibody
<b>Host</b>	FCGR1A antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Clone Name:</b>	10.1
<b>Conjugations:</b>	R. Phycoerythrin (RPE)
<b>Immunogen Species:</b>	FCGR1A / CD64 antibody was raised against Human
<b>Antigen Type:</b>	Cells
<b>Immunogen:</b>	FCGR1A / CD64 antibody was raised against human monocytes.
<b>Specificity:</b>	Recognizes the human CD64 cell surface antigen, a 75kD glycoprotein expressed by monocytes and by some myeloid cell lines and myeloid leukemias. CD64 is a high affinity receptor for IgG and is also known as FcRI
<b>Reactivity:</b>	Human, Baboon, Cynomolgus monkey, Rhesus monkey
<b>Purification:</b>	Protein G purified
<b>Reconstitution:</b>	Distilled Water.
<b>Presentation:</b>	Lyophilized, PBS, pH 7.4, 0.09% sodium azide, 1% BSA.
<b>Usage Summary:</b>	Flow Cytometry: Use 10 ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100 ul. Method sheets are available on request.
<b>Uses:</b>	Flow Cytometry (1:1 - 1:10) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	100 tst
<b>Requested From:</b>	Japan
<b>Laboratory Reagent For In Vitro Research Use Only</b>	
<b>Not for resale without prior written consent from LifeSpan BioSciences, Inc.</b>	
Created on 10/1/2014 © 2014 LifeSpan BioSciences	