

ICAM4 / CDE4 Mayoo a	oti Human Managlangi /FITC) (1H4) Antibody, LC C46409, LCBia
	nti-Human Monoclonal (FITC) (1H4) Antibody - LS-C46408 - LSBio
CatalogID:	LS-C46408
Target:	intercellular adhesion molecule 1 (ICAM1)
Synonyms:	ICAM1 Antibody, BB2 Antibody, CD54 antigen Antibody, CD54 Antibody, ICAM-1 Antibody, p3.58 Antibody
Family / Subfamily:	ICAM / not assigned-ICAM
Host	ICAM1 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG2b
Clone Name:	1H4
Conjugations:	Fluorescein (FITC)
Immunogen Species:	ICAM1 / CD54 antibody was raised against Human
Antigen Type:	Cells
Immunogen:	ICAM1 / CD54 antibody was raised against raji cells and spleen cells fused with NS1 cells.
Specificity:	Reacts with CD54 (ICAM-1), a 85-110 kD type I transmembrane glycoprotein (receptor for rhinovirus). The expression of CD54 is upregulated by activation; it is expressed on activated endothelial cells, T lymphocytes, B lymphocytes, monocytes, macrophages, granulocytes and dendritic cells.
Reactivity:	Human
Purification:	Purified
Presentation:	PBS, 15 mM sodium azide, 0.2% high-grade protease free BSA as a stabilizing agent.
Recommended Storage:	May be stored at 4°C before opening. DO NOT FREEZE! Stable at 4°C as an undiluted liquid. Dilute only prior to immediate use. Stable for at least 6 months at 4°C. Sensitive to light.
Usage Summary:	Use 20 ul per 100ul of whole blood or 10e6 cells for Flow cytometry.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 tst
Requested From:	Japan

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 10/1/2014

© 2014 LifeSpan BioSciences