

CatalogID:	LS-C4867
Target:	CD27 molecule
Synonyms:	CD27 Antibody, CD27 antigen Antibody, CD27 molecule Antibody, S152 Antibody, T14 Antibody, T-cell activation antigen CD27 Antibody, CD27L receptor Antibody, T cell activation antigen S152 Antibody, TNFRSF7 Antibody, Tp55 Antibody
Family / Subfamily:	TNF Receptor / not assigned-TNF Receptor
Host	CD27 antibody was produced in Mouse
Clonality:	Monoclonal
Isotype:	IgG2a
Conjugations:	Fluorescein (FITC)
Immunogen Species:	CD27 antibody was raised against Human
Specificity:	Precipitates a molecule of MW55kD (120kD in non-reduced conditions) designated as the CD27 antigen. This antigen is expressed on activated T lymphocytes and belongs to a family of proteins associated with the nerve growth factor receptor. CD27 is expressed by thymocytes and peripheral T cells and also by EBV transformed B cell lines. Species cross-reactivity: Cynomolgus monkey and Rhesus monkey.
Reactivity:	Human
Purification:	Ion exchange chromatography
Presentation:	Purified IgG.
Recommended Storage:	Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.
Usage Summary:	Flow Cytometry: Neat.
Uses:	Flow Cytometry (Optimal dilution to be determined by the researcher)
Size:	100 µg
Requested From:	Japan
La	boratory Reagent For In Vitro Research Use Only
Not for resale	without prior written consent from LifeSpan BioSciences, Inc.