

**Chlamydia trachomatis Goat anti-Chlamydia trachomatis Polyclonal (FITC) Antibody - LS-C50786 - LSBio**

<b>CatalogID:</b>	LS-C50786
<b>Target:</b>	Chlamydia trachomatis
<b>Host</b>	Goat
<b>Clonality:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Immunogen Species:</b>	Chlamydia trachomatis
<b>Immunogen:</b>	Chlamydia trachomatis, L2 + other serovar groups.
<b>Specificity:</b>	Purified elementary bodies, disrupted. Cross-reacts with Chlamydia psittacii & Chlamydia pneumoniae (TWAR). Uninfected Cell Reactivity: Negative vs. HEp-2 cells and egg yolk sac.
<b>Reactivity:</b>	Chlamydia trachomatis
<b>Purification:</b>	Purified
<b>Presentation:</b>	PBS, pH 7.2, BSA (10 mg/ml), 0.1% sodium azide.
<b>Recommended Storage:</b>	Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.
<b>Usage Summary:</b>	Suitable for use in Immunofluorescence. Immunofluorescence: 1:10-1:50. Direct FA staining of target antigens in a permissive tissue culture system. Acetone fixation of the antigen source is recommended prior to staining.
<b>Uses:</b>	Immunofluorescence (1:10 - 1:50) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	1,000 µl
<b>Requested From:</b>	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 10/2/2014

© 2014 LifeSpan BioSciences