

**Chlamydia trachomatis Rabbit anti-Chlamydia trachomatis Polyclonal (FITC) Antibody - LS-C50790  
 - LSBio**

|                             |  |
|-----------------------------|--|
| <b>CatalogID:</b>           | LS-C50790  |
| <b>Target:</b>              | Chlamydia trachomatis  |
| <b>Host</b>                 | Rabbit   |
| <b>Clonality:</b>           | Polyclonal   |
| <b>Isotype:</b>             | IgG  |
| <b>Conjugations:</b>        | Fluorescein (FITC)   |
| <b>Immunogen Species:</b>   | Chlamydia trachomatis  |
| <b>Immunogen:</b>           | Chlamydia trachomatis, L2 + other serovar groups.  |
| <b>Specificity:</b>         | Recognizes purified elementary bodies, disrupted. Species cross-reactivity: Chlamydia psittacii and Chlamydia pneumoniae (TWAR). Uninfected Cell Reactivity: Negative vs. HEp-2 cells and egg yolk sac.  |
| <b>Reactivity:</b>          | Chlamydia trachomatis  |
| <b>Purification:</b>        | Purified   |
| <b>Presentation:</b>        | PBS, pH 7.2, 10 mg/ml BSA, 0.1% sodium azide.  |
| <b>Recommended Storage:</b> | Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.   |
| <b>Usage Summary:</b>       | Suitable for use in Immunofluorescence. Immunofluorescence vs. all serovars (A-K, L1-L3): 1:10-1:50. Staining of target antigens in a permissive tissue culture system. Acetone fixation of the antigen source is recommended prior to staining. |
| <b>Uses:</b>                | Immunofluorescence (Optimal dilution to be determined by the researcher)   |
| <b>Size:</b>                | 1,000 µl   |
| <b>Requested From:</b>      | Japan  |

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 10/2/2014

© 2014 LifeSpan BioSciences