

**ICAM2 / CD102 Mouse anti-Human Monoclonal (PE) Antibody - LS-C5767 - LSBio**

<b>CatalogID:</b>	LS-C5767
<b>Target:</b>	intercellular adhesion molecule 2 (ICAM2)
<b>Synonyms:</b>	ICAM2 Antibody, CD102 Antibody, ICAM-2 Antibody, CD102 antigen Antibody
<b>Family / Subfamily:</b>	ICAM / not assigned-ICAM
<b>Host</b>	ICAM2 antibody was produced in Mouse
<b>Clonality:</b>	Monoclonal
<b>Isotype:</b>	IgG2a
<b>Conjugations:</b>	Phycoerythrin (PE)
<b>Immunogen Species:</b>	ICAM2 / CD102 antibody was raised against Human
<b>Antigen Type:</b>	Cells
<b>Immunogen:</b>	ICAM2 / CD102 antibody was raised against transfected COS cells expressing ICAM-2 cDNA.
<b>Specificity:</b>	Recognizes human CD102, also known as intercellular adhesion molecule-2 (ICAM-2).
<b>Reactivity:</b>	Human
<b>Purification:</b>	Protein G purified
<b>Reconstitution:</b>	deionized water. Possible additional volumes for resuspension: 200 µl
<b>Presentation:</b>	Lyophilized.
<b>Recommended Storage:</b>	Long term: Add glycerol (40-50%) -20°C; Short term: +4°C
<b>Usage Summary:</b>	Suitable for use in Flow Cytometry, ELISA, Immunoprecipitation. Flow Cytometry: 1:100. Use 10 ul of the suggested working dilution to label 1E6cells in 100ul. Immunohistochemistry: Frozen.
<b>Uses:</b>	IHC - Frozen, Immunoprecipitation, Flow Cytometry (1:100), ELISA (Optimal dilution to be determined by the researcher)
<b>Size:</b>	200 µg
<b>Requested From:</b>	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 10/2/2014

© 2014 LifeSpan BioSciences