

| CatalogID: | LS-C57998 |
|----------------------|--|
| Target: | CD28 molecule |
| Synonyms: | CD28 Antibody, CD28 antigen Antibody, CD28 antigen (Tp44) Antibody, Tp44 Antibody, CD28 molecule Antibody |
| Host | CD28 antibody was produced in Mouse |
| Clonality: | Monoclonal |
| Isotype: | IgG2b |
| Clone Name: | E18 |
| Conjugations: | R. Phycoerythrin (RPE) |
| Immunogen Species: | CD28 antibody was raised against Mouse |
| Immunogen: | CD28 antibody was raised against alternate immunization with A20 cells expressing mCD28 and recombinant mCD28lg. |
| Specificity: | Recognizes murine CD28, a co-stimulatory cell surface molecule expressed by T cell subsets. Has been shown to be able to stimulate murine T lymphocytes in combination with anti-mouse CD3 |
| Reactivity: | Mouse |
| Purification: | Affinity purified |
| Reconstitution: | Distilled Water. |
| Presentation: | Lyophilized, PBS, pH 7.4. |
| Recommended Storage: | +4°C, avoid freezing |
| Usage Summary: | Flow Cytometry: Use10 ul of the suggested working dilution to label 10^6 cells in 100 ul. Method sheets are available on request. |
| Uses: | Flow Cytometry (1:1 - 1:10) (Optimal dilution to be determined by the researcher) |
| Size: | 100 tst |
| Requested From: | Japan |
| La | boratory Reagent For In Vitro Research Use Only |
| Not for resale | without prior written consent from LifeSpan BioSciences, Inc. |