

**Bovine IgG Rabbit anti-Bovine Polyclonal (Fab'2) (PE) Antibody - LS-C61261 - LSBio**

<b>CatalogID:</b>	LS-C61261
<b>Target:</b>	Bovine IgG
<b>Host</b>	Bovine IgG antibody was produced in Rabbit
<b>Clonality:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Conjugations:</b>	Phycoerythrin (PE)
<b>Modifications:</b>	Fab'2
<b>Immunogen Species:</b>	Bovine IgG antibody was raised against Bovine
<b>Reactivity:</b>	Bovine
<b>Purification:</b>	Affinity purified
<b>Usage Summary:</b>	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain $1 \times 10^6$ cells in flow cytometry is approximately 1.0 ug of antibody conjugate. Lesser amounts of reagent may be sufficient for staining.
<b>Uses:</b>	Immunofluorescence (1:100 - 1:250), Flow Cytometry (1:100 - 1:250) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	500 µg
<b>Requested From:</b>	Japan

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 10/2/2014

© 2014 LifeSpan BioSciences