

## Mouse IgG Donkey anti-Mouse Polyclonal (Fab'2) (FITC) Antibody - LS-C61457 - LSBio

|                             |   |
|-----------------------------|---|
| <b>CatalogID:</b>           | LS-C61457   |
| <b>Target:</b>              | Mouse IgG   |
| <b>Host</b>                 | Mouse IgG antibody was produced in Donkey   |
| <b>Clonality:</b>           | Polyclonal  |
| <b>Isotype:</b>             | IgG   |
| <b>Conjugations:</b>        | Fluorescein (FITC)  |
| <b>Modifications:</b>       | Fab'2   |
| <b>Immunogen Species:</b>   | Mouse IgG antibody was raised against Mouse   |
| <b>Antigen Type:</b>        | Purified protein  |
| <b>Immunogen:</b>           | Mouse IgG antibody was raised against mouse IgG whole molecule.   |
| <b>Specificity:</b>         | anti-Fluorescein, anti-Donkey Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Sheep Serum Proteins                  |
| <b>Reactivity:</b>          | Mouse   |
| <b>Purification:</b>        | Immunoaffinity purified   |
| <b>Presentation:</b>        | 0.01 M sodium phosphate, 0.25 M sodium chloride, pH 7.2, 10 mg/ml BSA (IgG, protease free), 0.01% sodium azide.   |
| <b>Recommended Storage:</b> | Store at 4°C.   |
| <b>Usage Summary:</b>       | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |
| <b>Uses:</b>                | Immunofluorescence (1:50 - 1:200), Flow Cytometry (1:50 - 1:200) (Optimal dilution to be determined by the researcher)  |
| <b>Size:</b>                | 500 µg  |
| <b>Requested From:</b>      | Japan   |

Laboratory Reagent For In Vitro Research Use Only

Not for resale without prior written consent from LifeSpan BioSciences, Inc.

Created on 10/2/2014

© 2014 LifeSpan BioSciences