

Rabbit IgG Fab'2 Goat anti-Rabbit Polyclonal (Fab'2) (FITC) Antibody - LS-C61476 - LSBio	
<b>CatalogID:</b>	LS-C61476
<b>Target:</b>	Rabbit IgG Fab'2
<b>Host</b>	Goat
<b>Clonality:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Conjugations:</b>	Fluorescein (FITC)
<b>Modifications:</b>	Fab'2
<b>Immunogen Species:</b>	Rabbit
<b>Immunogen:</b>	Rabbit IgG F(ab') <sub>2</sub> fragment.
<b>Specificity:</b>	anti-Fluorescein, anti- Goat Serum, Rabbit IgG, Rabbit IgG F(ab') <sub>2</sub> and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rabbit IgG F(c) or Bovine, Horse, Human, Mouse, Rat and Sheep Serum Proteins
<b>Reactivity:</b>	Rabbit
<b>Purification:</b>	Immunoaffinity purified
<b>Reconstitution:</b>	deionized water. Possible additional volumes for resuspension: 500 µl
<b>Presentation:</b>	Lyophilized, 0.02 M potassium phosphate, 0.15 M sodium chloride, pH 7.2, 10 mg/ml BSA (IgG, protease free), 0.01% sodium azide.
<b>Recommended Storage:</b>	Short term +4C. Aliquot to avoid freeze/thaw cycles and store at -20C. FITC conjugates are sensitive to light.
<b>Usage Summary:</b>	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.
<b>Uses:</b>	Immunofluorescence (1:500 - 1:2500), Flow Cytometry (1:2000 - 1:10000) (Optimal dilution to be determined by the researcher)
<b>Size:</b>	500 µg
<b>Requested From:</b>	Japan
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioSciences, Inc.	
Created on 10/2/2014	
© 2014 LifeSpan BioSciences	