

antigen Antibody, Leu2 T-lymphocyte antigen Antibody, T8 T-cell antigen Antibody, manifer Cell Co-receptor Antibody, p32 Antibody, CD8 Antibody, Leu2 Antibody, T-cell antigen Leu2 Antibody Host CD8A antibody was produced in Rat Clonality: Monoclonal Isotype: IgG2a Clone Name: CT-CD8a Conjugations: Fluorescein (FITC) Immunogen Species: CD8A / CD8 Alpha antibody was raised against Mouse Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Purified Presentation: PBS, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be deterr by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan	CD8A Antibody, CD8a molecule Antibody, CD8a antigen Antibody, OKT8 T-cell antigen Antibody, Leu2 T-lymphocyte antigen Antibody, T8 T-cell antigen Antibody T cell co-receptor Antibody, p32 Antibody, CD8 Antibody, Leu2 Antibody, T-cell antigen Leu2 Antibody
antigen Antibody, Leu2 T-lymphocyte antigen Antibody, TB T-cell antigen Antibody, p32 Antibody, CD8 Antibody, Leu2 Antibody, T-cell antigen Leu2 Antibody Host CD8A antibody was produced in Rat Clonality: Monoclonal Isotype: IgG2a Clone Name: CT-CD8a Conjugations: Fluorescein (FITC) Immunogen Species: CD8A / CD8 Alpha antibody was raised against Mouse Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Pus, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be deterr by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan	antigen Antibody, Leu2 T-lymphocyte antigen Antibody, T8 T-cell antigen Antibody T cell co-receptor Antibody, p32 Antibody, CD8 Antibody, Leu2 Antibody, T-cell antigen Leu2 Antibody
Clonality: Monoclonal Isotype: IgG2a Clone Name: CT-CD8a Conjugations: Fluorescein (FITC) Immunogen Species: CD8A / CD8 Alpha antibody was raised against Mouse Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Purified Presentation: PBS, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be deterr by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	
Isotype: IgG2a Clone Name: CT-CD8a Conjugations: Fluorescein (FITC) Immunogen Species: CD8A / CD8 Alpha antibody was raised against Mouse Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Purified Presentation: PBS, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be deterr by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	CD8A antibody was produced in Rat
Clone Name: CT-CD8a Conjugations: Fluorescein (FITC) Immunogen Species: CD8A / CD8 Alpha antibody was raised against Mouse Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Purified Presentation: PBS, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be deterr by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Monoclonal
Conjugations: Fluorescein (FITC) Immunogen Species: CD8A / CD8 Alpha antibody was raised against Mouse Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Purified Presentation: PBS, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be determ by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	IgG2a
Immunogen Species: CD8A / CD8 Alpha antibody was raised against Mouse Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Purified Presentation: PBS, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be determ by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	CT-CD8a
Specificity: Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 a chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict. Reactivity: Mouse Purification: Purified Presentation: PBS, 0.1% sodium azide, EIA grade BSA. Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be determ by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Fluorescein (FITC)
chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8alpha/beta heterodimer that is expressed by the majority of thymocytes a the MHC class I restrict.Reactivity:MousePurification:PurifiedPresentation:PBS, 0.1% sodium azide, EIA grade BSA.Recommended Storage:Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycleUses:IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be deterr by the researcher)Size:100 µg or 300 µgConcentration:0.1 mg/mlRequested From:JapanLaboratory Reagent For In Vitro Research Use Only	CD8A / CD8 Alpha antibody was raised against Mouse
Purification:PurifiedPresentation:PBS, 0.1% sodium azide, EIA grade BSA.Recommended Storage:Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycleUses:IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be determ by the researcher)Size:100 µg or 300 µgConcentration:0.1 mg/mlRequested From:JapanLaboratory Reagent For In Vitro Research Use Only	CD8alpha/beta heterodimer that is expressed by the majority of thymocytes and by
Presentation:PBS, 0.1% sodium azide, EIA grade BSA.Recommended Storage:Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycleUses:IHC - Frozen, Flow Cytometry (1 μg/10E6 cells) (Optimal dilution to be deterr by the researcher)Size:100 μg or 300 μgConcentration:0.1 mg/mlRequested From:JapanLaboratory Reagent For In Vitro Research Use Only	Mouse
Recommended Storage: Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycle Uses: IHC - Frozen, Flow Cytometry (1 µg/10E6 cells) (Optimal dilution to be determed by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Purified
Uses: IHC - Frozen, Flow Cytometry (1 μg/10E6 cells) (Optimal dilution to be deterr by the researcher) Size: 100 μg or 300 μg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	PBS, 0.1% sodium azide, EIA grade BSA.
by the researcher) by the researcher) Size: 100 µg or 300 µg Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.
Concentration: 0.1 mg/ml Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	IHC - Frozen, Flow Cytometry (1 μg /10E6 cells) (Optimal dilution to be determined by the researcher)
Requested From: Japan Laboratory Reagent For In Vitro Research Use Only	100 µg or 300 µg
Laboratory Reagent For In Vitro Research Use Only	0.1 mg/ml
· · · · · · · · · · · · · · · · · · ·	Japan
	ooratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc. Created on 10/2/2014	
Not for resale v	