

| CD8A / CD8 Alpha Rat anti-Mouse Monoclonal (PE, Cy5) (CT-CD8a) Antibody - LS-C62784 - LSBio |  |
|---|--|
| <b>CatalogID:</b>   | LS-C62784  |
| <b>Target:</b>  | CD8a molecule  |
| <b>Synonyms:</b>  | CD8A Antibody, CD8a molecule Antibody, CD8a antigen Antibody, OKT8 T-cell antigen Antibody, Leu2 T-lymphocyte antigen Antibody, T8 T-cell antigen Antibody, T cell co-receptor Antibody, p32 Antibody, CD8 Antibody, Leu2 Antibody, T-cell antigen Leu2 Antibody |
| <b>Host</b>   | CD8A antibody was produced in Rat  |
| <b>Clonality:</b>   | Monoclonal   |
| <b>Isotype:</b>   | IgG2a  |
| <b>Clone Name:</b>  | CT-CD8a  |
| <b>Conjugations:</b>  | Phycoerythrin (PE), Cy5  |
| <b>Immunogen Species:</b>   | CD8A / CD8 Alpha antibody was raised against Mouse   |
| <b>Specificity:</b>   | Anti-mouse CD8a antigen monoclonal antibody recognizes the mouse CD8 alpha chain. The alpha chain of CD8 associates with the CD8 beta chain to form a CD8 alpha/beta heterodimer that is expressed by the majority of thymocytes and by the MHC class I restric. |
| <b>Reactivity:</b>  | Mouse  |
| <b>Purification:</b>  | Purified   |
| <b>Presentation:</b>  | PBS, 0.1% sodium azide, EIA grade BSA.   |
| <b>Uses:</b>  | Flow Cytometry (1 µg/10E6 cells), Cytotoxic Assay (Optimal dilution to be determined by the researcher)  |
| <b>Size:</b>  | 100 µg   |
| <b>Requested From:</b>  | Japan  |
| <b>Laboratory Reagent For In Vitro Research Use Only</b>                                    |  |
| Not for resale without prior written consent from LifeSpan BioSciences, Inc.                |  |
| Created on 10/2/2014  |  |
| © 2014 LifeSpan BioSciences   |  |