

Clonality: Monoclonal Isotype: IgG1 Clone Name: 9 E10 Immunogen Species: MYC / c-Myc antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: MYC / c-Myc antibody was raised against synthetic peptide residues 408-439 of the human p62c-myc protein. Specificity: Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody cross reacts with both commyc-p64c-myc protein. Epitope: aa408-439 Reactivity: Human, Mouse Purification: Ascites Presentation: 0.05% sodium azide Recommended Storage: Store at 2°C to 8°C degrees. Do not freeze. Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)	
oncogene protein Antibody, Proto-oncogene c-Myc Antibody Transcription factor p64 Antibody, v-myc Antibody Host MYC antibody was produced in Mouse Clonality: Monoclonal Isotype: IgG1 Clone Name: 9 E10 Immunogen Species: MYC / c-Myc antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: MYC / c-Myc antibody was raised against synthetic peptide residues 408-439 of the human p62c-myc protein. Specificity: Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody cross reacts with human p62/6 react with mouse p62/64c-myc protein. Epitope: aa408-439 Reactivity: Human, Mouse Purification: Ascites Presentation: 0.05% sodium azide Recommended Storage: Store at 2°C to 8°C degrees. Do not freeze. Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher) Size: 0.5 ml or 1 ml Requested From: Japan	
Clonality: Monoclonal Isotype: IgG1 Clone Name: 9 E10 Immunogen Species: MYC / c-Myc antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: MYC / c-Myc antibody was raised against synthetic peptide of residues 408-439 of the human p62c-myc protein. Specificity: Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody cross reacts with human p62/c react with mouse p62/64c-myc protein. Epitope: aa408-439 Reactivity: Human, Mouse Purification: Ascites Presentation: 0.05% sodium azide Recommended Storage: Store at 2°C to 8°C degrees. Do not freeze. Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher) Size: 0.5 ml or 1 ml Requested From: Japan	Antibody, Myc proto- y, Rats1 Antibody,
Isotype: IgG1 Clone Name: 9 E10 Immunogen Species: MYC / c-Myc antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: MYC / c-Myc antibody was raised against synthetic peptide residues 408-439 of the human p62c-myc protein. Specificity: Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody cross reacts with human p62/6 react with mouse p62/64c-myc protein. Epitope: aa408-439 Reactivity: Human, Mouse Purification: Ascites Presentation: 0.05% sodium azide Recommended Storage: Store at 2°C to 8°C degrees. Do not freeze. Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher) Size: 0.5 ml or 1 ml Requested From: Japan	
Clone Name: 9 E10 Immunogen Species: MYC / c-Myc antibody was raised against Human Antigen Type: Synthetic peptide Immunogen: MYC / c-Myc antibody was raised against synthetic peptide residues 408-439 of the human p62c-myc protein. Specificity: Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody reacts with human p62/c react with mouse p62/64c-myc protein. Epitope: aa408-439 Reactivity: Human, Mouse Purification: Ascites Presentation: 0.05% sodium azide Recommended Storage: Store at 2°C to 8°C degrees. Do not freeze. Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher) Size: 0.5 ml or 1 ml Requested From: Japan	
Immunogen Species:MYC / c-Myc antibody was raised against HumanAntigen Type:Synthetic peptideImmunogen:MYC / c-Myc antibody was raised against synthetic peptide or residues 408-439 of the human p62c-myc protein.Specificity:Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody cross reacts with human p62cEpitope:aa408-439Reactivity:Human, MousePurification:AscitesPresentation:0.05% sodium azideRecommended Storage:Store at 2°C to 8°C degrees. Do not freeze.Uses:Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)Size:0.5 ml or 1 mlRequested From:Japan	
Antigen Type: Synthetic peptide Immunogen: MYC / c-Myc antibody was raised against synthetic peptide or residues 408-439 of the human p62c-myc protein. Specificity: Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both commyc-p64c-myc. This antibody cross reacts with human p62/6 react with mouse p62/64c-myc protein. Epitope: aa408-439 Reactivity: Human, Mouse Purification: Ascites Presentation: 0.05% sodium azide Recommended Storage: Store at 2°C to 8°C degrees. Do not freeze. Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher) Size: 0.5 ml or 1 ml Requested From: Japan	
Immunogen: MYC / c-Myc antibody was raised against synthetic peptide of residues 408-439 of the human p62c-myc protein. Specificity: Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody cross reacts with human p62/6 react with mouse p62/64c-myc protein. Epitope: aa408-439 Reactivity: Human, Mouse Purification: Ascites Presentation: 0.05% sodium azide Recommended Storage: Store at 2°C to 8°C degrees. Do not freeze. Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher) Size: 0.5 ml or 1 ml Requested From: Japan	
residues 408-439 of the human p62c-myc protein.Specificity:Recognizes an epitope located in the amino acid residue 41 oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody cross reacts with human p62/6 react with mouse p62/64c-myc protein.Epitope:aa408-439Reactivity:Human, MousePurification:AscitesPresentation:0.05% sodium azideRecommended Storage:Store at 2°C to 8°C degrees. Do not freeze.Uses:Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)Size:0.5 ml or 1 mlRequested From:Japan	
oncogene product c-myc. This antibody reacts with both con myc-p64c-myc. This antibody cross reacts with human p62/6 react with mouse p62/64c-myc protein.Epitope:aa408-439Reactivity:Human, MousePurification:AscitesPresentation:0.05% sodium azideRecommended Storage:Store at 2°C to 8°C degrees. Do not freeze.Uses:Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)Size:0.5 ml or 1 mlRequested From:Japan	corresponding to
Reactivity:Human, MousePurification:AscitesPresentation:0.05% sodium azideRecommended Storage:Store at 2°C to 8°C degrees. Do not freeze.Uses:Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)Size:0.5 ml or 1 mlRequested From:Japan	mponents of the p62c-
Purification:AscitesPresentation:0.05% sodium azideRecommended Storage:Store at 2°C to 8°C degrees. Do not freeze.Uses:Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)Size:0.5 ml or 1 mlRequested From:Japan	
Presentation:0.05% sodium azideRecommended Storage:Store at 2°C to 8°C degrees. Do not freeze.Uses:Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)Size:0.5 ml or 1 mlRequested From:Japan	
Recommended Storage:Store at 2°C to 8°C degrees. Do not freeze.Uses:Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher)Size:0.5 ml or 1 mlRequested From:Japan	
Uses: Western blot (1:100 - 1:500), Immunoprecipitation (Optimal of determined by the researcher) Size: 0.5 ml or 1 ml Requested From: Japan	
determined by the researcher) Size: 0.5 ml or 1 ml Requested From:	
Requested From: Japan	dilution to be
Laboratory Reagent For In Vitro Research Use Only	
Not for resale without prior written consent from LifeSpan BioScience Created on 10/2/2014	es, Inc.