

Product Datasheet

Mas Antibody NBP1-7844SS

Unit Size: 0.025 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



support@novusbio.com

Reviews: 2 Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-78444

Updated 6/15/2014 v.20.1

NBP1-78444SS


Mas Antibody

Product Information	
Unit Size	0.025 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Purity	Immunogen affinity purified
Buffer	PBS, 30% glycerol
Product Description	
Host	Rabbit
Gene ID	4142
Gene Symbol	MAS1
Species	Human, Mouse, Rat, Primate, Rabbit
Species Reactivity	Human, monkey, rat, and mouse. Rabbit reactivity reported in the scientific literature (PMID: 23701246).
Immunogen	A synthetic peptide made to an internal portion of the human MAS1 protein (between residues 75-125) [UniProt P04201]
Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunocytochemistry/Immunofluorescence 1:25, Immunohistochemistry 1:100, Immunohistochemistry-Paraffin 1:100, Western Blot 1:1000, Simple Western 1:100
Application Notes	This MAS1 antibody is useful for Immunocytochemistry/Immunofluorescence, Immunohistochemistry on paraffin embedded sections and Western blot. In Simple Western only 10-15 uL of the recommended dilution is used per data point.




Images


Western Blot: MAS1 Antibody [NBP1-78444] - Analysis of MAS1 in: 1. Ntera 2, 2. A431, 3. HepG2, 4. MCF7, 5. 3T3 and 6. Cos7.



Immunocytochemistry/Immunofluorescence: MAS1 Antibody [NBP1-78444] - MAS1 antibody was tested at 1:25 in HepG2 cells with FITC (green). Nuclei were counterstained with Dapi (blue).



Immunohistochemistry: MAS1 Antibody [NBP1-78444] - Analysis of MAS1 in mouse kidney using DAB with hematoxylin counterstain.



Simple Western: Mas Antibody [NBP1-78444] - Simple Western lane view shows a specific band for MAS1 in 0.5 mg/ml of HepG2 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Publications

Bian W, Sun L, Yang L et al. Stabilization of the Angiotensin-(1-7) Receptor Mas through Interaction with PSD95. *Biochem J* 2013 May 24 [PMID: 23701246] (WB, Rabbit)

da Silveira KD, Coelho FM, Vieira AT et al. Anti-inflammatory effects of the activation of the angiotensin-(1-7) receptor, MAS, in experimental models of arthritis *J Immunol* 2010 Nov 1 [PMID: 20935211] (WB, Mouse)



Procedures

Western Blot protocol for MAS1 Antibody (NBP1-78444)

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
 2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
 3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
 4. Rinse the blot.
 5. Block the membrane using standard blocking buffer for at least 1 hour.
 6. Wash the membrane in wash buffer three times for 10 minutes each.
 7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
 8. Wash the membrane in wash buffer three times for 10 minutes each.
 9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
 10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
 11. Apply the detection reagent of choice in accordance with the manufacturers instructions.
- Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

Immunohistochemistry-Paraffin protocol for MAS1 Antibody (NBP1-78444)

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



Immunocytochemistry/Immunofluorescence Protocol for MAS1 Antibody (NBP1-78444)

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





Novus Biologicals USA

8100 Southpark Way, A-8
Littleton, CO 80120
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Novus Biologicals Canada

461 North Service Road West, Unit B37
Oakville, ON L6M 2V5
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada@novusbio.com

Novus Biologicals Europe

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our guarantee, please visit www.novusbio.com/guarantee.

