# **Product Datasheet**

## Bladder Lysate NBP2-29015

Unit Size: 3 Vials

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



support@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-29015

Updated 1/9/2014 v.20.1

### NBP2-29015

Human Bladder Tissue Lysate (Adult)

| Human Diadder Hissue Lysate (Addit) |  |
|-------------------------------------|--|
| Product Information                 |  |
| Unit Size                           | 3 Vials  |
| Concentration                       | 1 mg/ml  |
| Storage                             | Store at -80C. Avoid freeze-thaw cycles.   |
| Reconstitution Instructions         | The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides.   |
| Product Description                 |  |
| Species                             | Human  |
| Specificity/Sensitivity             | Age :64<br>Diagnosis:Transitional cell carcinoma. Moderately differentiated<br>Grade:2<br>Stage:II<br>TNM:T2N1M0   |
| Immunogen                           | Clinical Tissue<br>Human Bladder Tissue Lysate Bundle Set  |
| Preparation Method                  | Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. |
| Lysate Type                         | Tissue   |
| Lysate Tissue                       | Bladder  |
| Lysate Life Stage                   | Adult  |
| Product Application Details         |  |
| Application Notes                   | These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required.                          |





#### **Novus Biologicals USA**

8100 Southpark Way, A-8 Littleton, CO 80120 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 novus@novusbio.com

#### **Novus Biologicals Canada**

461 North Service Road West, Unit B37 Oakville, ON L6M 2V5 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada@novusbio.com

#### **Novus Biologicals Europe**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: technical@novusbio.com Orders: orders@novusbio.com General: novus@novusbio.com

#### Products Related to NBP2-29015

NBP2-30252

Bladder Tissue Slides (Transitional Cell Carcinoma)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our guarantee, please visit www.novusbio.com/guarantee.

