

# **Product Datasheet**

## **Kappa Light Chain Antibody**

### **NBP2-33142-0.2mg**

Unit Size: 0.2 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[support@novusbio.com](mailto:support@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-33142](http://www.novusbio.com/NBP2-33142)

Updated 11/13/2014 v.20.1

**NBP2-33142-0.2mg**

Kappa Light Chain Antibody (L1C1) [Azide Free]

<b>Product Information</b>	
<b>Unit Size</b>	0.2 mg
<b>Concentration</b>	Please see the vial label for concentration. If unlisted please contact technical services.
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	L1C1
<b>Preservative</b>	No Preservative
<b>Isotype</b>	IgG1 Kappa
<b>Conjugate</b>	Azide Free
<b>Purity</b>	Protein G purified
<b>Buffer</b>	10mM PBS. Does not contain BSA.

<b>Product Description</b>	
<b>Host</b>	Mouse
<b>Gene ID</b>	3514
<b>Gene Symbol</b>	IGKC
<b>Species</b>	Human
<b>Specificity/Sensitivity</b>	This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain, kappa or lambda. The ratio of kappa to lambda is 70:30. However, with the occurrence of multiple myeloma or other B-cell malignancies this ratio is disturbed. Antibody to the kappa light chain is reportedly useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkin's lymphomas. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is malignant.
<b>Immunogen</b>	B lymphoma cells

<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, ELISA, Flow Cytometry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation
<b>Recommended Dilutions</b>	Flow Cytometry 0.5-1ug/million cells, Immunoprecipitation 1-2ug/500ug protein lysate, Western Blot 1:250-1:500, Immunohistochemistry-Frozen 0.5-1.0ug/ml, Immunohistochemistry-Paraffin 0.5-1.0ug/ml, ELISA 1-5ug/ml for coating
<b>Application Notes</b>	Immunohistochemical staining of formalin/paraffin tissues is enhanced by unmasking of the target epitope by boiling the tissue sections in 10mM citrate buffer, pH 6.0, for 10 min followed by cooling at room temp for 20 min.





### **Novus Biologicals USA**

8100 Southpark Way, A-8  
Littleton, CO 80120  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
novus@novusbio.com

### **Novus Biologicals Canada**

461 North Service Road West, Unit B37  
Oakville, ON L6M 2V5  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada@novusbio.com

### **Novus Biologicals Europe**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: technical@novusbio.com  
Orders: orders@novusbio.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

**For more information on our guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee).**

