

# Product Datasheet

## Cytokeratin 8 Antibody NBP2-34501-0.2 mg

Unit Size: 0.2 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[support@novusbio.com](mailto:support@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-34501](http://www.novusbio.com/NBP2-34501)

Updated 7/6/2014 v.20.1

**NBP2-34501-0.2 mg**

Cytokeratin 8 Antibody (TS1) [Azide Free]

| <b>Product Information</b>         |   |
|------------------------------------|---|
| <b>Unit Size</b>                   | 0.2 mg  |
| <b>Concentration</b>               | Please see the vial label for concentration. If unlisted please contact technical services.   |
| <b>Storage</b>                     | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.  |
| <b>Clonality</b>                   | Monoclonal  |
| <b>Clone</b>                       | TS1   |
| <b>Preservative</b>                | No Preservative   |
| <b>Isotype</b>                     | IgG1  |
| <b>Conjugate</b>                   | Azide Free  |
| <b>Purity</b>                      | Protein A purified  |
| <b>Buffer</b>                      | 10 mM PBS. Does not contain BSA.  |
| <b>Target Molecular Weight</b>     | 52.5 kDa  |
| <b>Product Description</b>         |   |
| <b>Host</b>                        | Mouse   |
| <b>Gene ID</b>                     | 3856  |
| <b>Gene Symbol</b>                 | KRT8  |
| <b>Species</b>                     | Human, Rat (Negative)   |
| <b>Immunogen</b>                   | Keratin preparation from a human carcinoma  |
| <b>Product Application Details</b> |   |
| <b>Applications</b>                | Western Blot, ELISA, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation  |
| <b>Recommended Dilutions</b>       | Flow Cytometry 0.5-1ug/million cells, Immunocytochemistry/Immunofluorescence 1-2ug/ml, Immunoprecipitation 1-2ug/500ug protein, Western Blot 0.5-1.0ug/ml, Immunohistochemistry-Frozen 0.5-1.0ug/ml, Immunohistochemistry-Paraffin 0.5-1.0ug/ml, ELISA 1-5ug/ml for coating |
| <b>Application Notes</b>           | Immunohistochemistry (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)   |





### **Novus Biologicals USA**

8100 Southpark Way, A-8  
Littleton, CO 80120  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
novus@novusbio.com

### **Novus Biologicals Canada**

461 North Service Road West, Unit B37  
Oakville, ON L6M 2V5  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada@novusbio.com

### **Novus Biologicals Europe**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: technical@novusbio.com  
Orders: orders@novusbio.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

**For more information on our guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee).**

