

# Product Datasheet

## **CD34 Antibody** **NBP2-34733-0.2 mg**

Unit Size: 0.2 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[support@novusbio.com](mailto:support@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-34733](http://www.novusbio.com/NBP2-34733)

Updated 7/6/2014 v.20.1

**NBP2-34733-0.2 mg**

CD34 Antibody (SPM123) [Azide Free]

<b>Product Information</b>	
<b>Unit Size</b>	0.2 mg
<b>Concentration</b>	Please see the vial label for concentration. If unlisted please contact technical services.
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	SPM123
<b>Preservative</b>	No Preservative
<b>Isotype</b>	IgG1 Kappa
<b>Conjugate</b>	Azide Free
<b>Purity</b>	Protein A purified
<b>Buffer</b>	10 mM PBS. Does not contain BSA
<b>Target Molecular Weight</b>	100 kDa
<b>Product Description</b>	
<b>Host</b>	Mouse
<b>Gene ID</b>	947
<b>Gene Symbol</b>	CD34
<b>Species</b>	Human, Primate, Bovine (Negative), Canine (Negative), Rat (Negative), Sheep (Negative)
<b>Species Reactivity</b>	Cynomolgus Monkey, Rhesus Monkey
<b>Specificity/Sensitivity</b>	This MAb recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34.
<b>Immunogen</b>	Detergent solubilized vesicular suspension prepared from human term placenta
<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation
<b>Recommended Dilutions</b>	Flow cytometry 0.5-1ug/million cells, Immunocytochemistry/Immunofluorescence 0.5-1ug/ml, Western Blot 0.25-0.5ug/ml, Immunoprecipitation 0.5-1ug/500ug protein lysate, Immunohistochemistry-Frozen 0.5-1ug/ml, immunohistochemistry-Paraffin 0.5-1ug/ml
<b>Application Notes</b>	Immunohistochemistry (Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)





### **Novus Biologicals USA**

8100 Southpark Way, A-8  
Littleton, CO 80120  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
novus@novusbio.com

### **Novus Biologicals Canada**

461 North Service Road West, Unit B37  
Oakville, ON L6M 2V5  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada@novusbio.com

### **Novus Biologicals Europe**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: technical@novusbio.com  
Orders: orders@novusbio.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

**For more information on our guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee).**

