# **Product Datasheet**

## CD63 Antibody NBP2-34779-0.2 mg

Unit Size: 0.2 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



support@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34779

Updated 7/6/2014 v.20.1

### NBP2-34779-0.2 mg


CD63 Antibody (SPM524) [Azide Free]

| Product Information         |   |
|-----------------------------|---|
| Unit Size                   | 0.2 mg  |
| Concentration               | Please see the vial label for concentration. If unlisted please contact technical services.   |
| Storage                     | Store at -20 to -80C. Avoid freeze-thaw cycles.   |
| Clonality                   | Monoclonal  |
| Clone                       | SPM524  |
| Preservative                | No Preservative   |
| Isotype                     | IgG1 Kappa  |
| Conjugate                   | Azide Free  |
| Purity                      | Protein A purified  |
| Buffer                      | 10 mM PBS. Does not contain BSA   |
| Product Description         |   |
| Host                        | Mouse   |
| Gene ID                     | 967   |
| Gene Symbol                 | CD63  |
| Species                     | Human, Mouse  |
| Specificity/Sensitivity     | The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-<br>molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression. |
| Immunogen                   | Smooth plasma membrane fraction of MeWo cells   |
| Product Application Details |   |
| Applications                | Western Blot, ELISA, Immunocytochemistry/Immunofluorescence,<br>Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin,<br>Immunoprecipitation  |
| Recommended Dilutions       | Immunocytochemistry/Immunofluorescence 0.5-1ug/ml, Western Blot 0.5-1ug/ml<br>Immunoprecipitation 0.5-1ug/500ug protein lysate, Immunohistochemistry-<br>Frozen 0.5-1ug/ml, Immunohistochemistry-Paraffin 0.5-1ug/ml, ELISA 1-5ug/ml<br>for coating   |
| Application Notes           | Immunohistochemistry (Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)   |



#### Images

Immunohistochemistry-Paraffin: CD63 Antibody (SPM524) [Azide Free] [NBP2-34779] - Formalin-paraffin human melanoma stained with CD63 MAb (SPM524)



Immunohistochemistry-Paraffin: CD63 Antibody (SPM524) [Azide Free] [NBP2-34779] - Formalin-paraffin mouse spleen stained with CD63 MAb (SPM524)







#### **Novus Biologicals USA**

8100 Southpark Way, A-8 Littleton, CO 80120 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 novus@novusbio.com

#### **Novus Biologicals Canada**

461 North Service Road West, Unit B37 Oakville, ON L6M 2V5 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada@novusbio.com

#### **Novus Biologicals Europe**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: technical@novusbio.com Orders: orders@novusbio.com General: novus@novusbio.com

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our guarantee, please visit www.novusbio.com/guarantee.

