Product Datasheet

Vimentin Antibody NBP2-34809-0.2 mg

Unit Size: 0.2 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



support@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34809

Updated 7/6/2014 v.20.1

NBP2-34809-0.2 mg

Vimentin Antibody (SPM576) [Azide Free]

| Product Information | |
|-----------------------------|--|
| Unit Size | 0.2 mg |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. |
| Clonality | Monoclonal |
| Clone | SPM576 |
| Preservative | No Preservative |
| Isotype | IgG1 |
| Conjugate | Azide Free |
| Purity | Protein A purified |
| Buffer | 10 mM PBS. Does not contain BSA |
| Target Molecular Weight | 58 kDa |
| Product Description | |
| Host | Mouse |
| Gene ID | 7431 |
| Gene Symbol | VIM |
| Species | Human, Bovine, Canine, Chicken, Feline, Goat, Porcine, Mouse (Negative), Rat (Negative) |
| Specificity/Sensitivity | This MAb reacts with a 58kDa protein identified as vimentin. It shows no cross- reaction with other closely related intermediate filament proteins (IFP's) such as desmin, keratin, neurofilament, and glial fibrillary acid protein. |
| Immunogen | Human vimentin recombinant protein |
| Product Application Details | |
| Applications | Western Blot, ELISA, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation |
| Recommended Dilutions | Flow Cytometry 0.5-1ug/million cells, Immunocytochemistry/Immunofluorescence 1-2ug/ml, Western Blot 0.5-1ug/ml, Immunoprecipitation 1-2ug/500ug protein lysate, Immunohistochemistry-Frozen 0.5-1ug/ml, Immunohistochemistry-Paraffin 0.5-1ug/ml, ELISA 1-5ug/ml for coating |
| Application Notes | Immunohistochemistry (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes) |





Novus Biologicals USA

8100 Southpark Way, A-8 Littleton, CO 80120 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 novus@novusbio.com

Novus Biologicals Canada

461 North Service Road West, Unit B37 Oakville, ON L6M 2V5 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada@novusbio.com

Novus Biologicals Europe

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: technical@novusbio.com Orders: orders@novusbio.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our guarantee, please visit www.novusbio.com/guarantee.

