

# Product Datasheet

## FPRL1/FPR2 Antibody NLS1878SS

Unit Size: 0.025 ml

Store at 4C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[support@novusbio.com](mailto:support@novusbio.com)

**Reviews: 1 Publications: 3**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NLS1878](http://www.novusbio.com/NLS1878)

Updated 6/15/2014 v.20.1

**NLS1878SS**

FPRL1/FPR2 Antibody

<b>Product Information</b>	
<b>Unit Size</b>	0.025 ml
<b>Concentration</b>	1.1 mg/ml
<b>Storage</b>	Store at 4C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Polyclonal
<b>Preservative</b>	0.01% Sodium Azide
<b>Purity</b>	Immunogen affinity purified
<b>Buffer</b>	PBS, 30% glycerol
<b>Target Molecular Weight</b>	38 kDa
<b>Product Description</b>	
<b>Host</b>	Rabbit
<b>Gene ID</b>	2358
<b>Gene Symbol</b>	FPR2
<b>Species</b>	Human, Mouse, Rat
<b>Species Reactivity</b>	Human and mouse. Rat reactivity reported in scientific literature (PMID: 24086560)
<b>Immunogen</b>	Synthetic peptide made to the 2nd extracellular loop of the human FPRL1 protein (between residues 300-350) [UniProt P25090]
<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
<b>Recommended Dilutions</b>	Immunocytochemistry/Immunofluorescence 1:20-1:75, Immunohistochemistry 1:300, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin 1:300, Western Blot 1:1000
<b>Application Notes</b>	This FPRL1 antibody is useful for Western Blot, Immunocytochemistry/Immunofluorescence, and Immunohistochemistry paraffin embedded sections. In Western Blot, a band is seen ~38 kDa representing FPRL1. In ICC/IF, plasma membrane staining was observed in Raw264.7 cells. In IHC-P, staining was also observed in the plasma membrane of human kidney cancer tissue. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended. Customers have reported success in IF on FFPE mouse kidney tissue, following microwave antigen retrieval. Use in Immunohistochemistry-Frozen reported in scientific literature (PMID 24086560)




## Images


Western Blot: FPRL1 Antibody [NLS1878] - WB analysis of FPRL1 in HL-60 cell lysate.



Immunocytochemistry/Immunofluorescence: FPRL1 Antibody [NLS1878] - FPRL1 antibody was tested in Raw264.7 cells with Dylight 488 (green). Nuclei were counterstained with DAPI (blue).



Immunohistochemistry: FPRL1 Antibody [NLS1878] - IHC analysis of FPRL1 in human kidney cancer using DAB with hematoxylin counterstain.



## Publications

Abdelmoaty S, Wigerblad G, Bas DB et al. Spinal Actions of Lipoxin A4 and 17(R)-Resolvin D1 Attenuate Inflammation-Induced Mechanical Hypersensitivity and Spinal TNF Release. PLoS One. 2013 Sep 24 [PMID: 24086560] (IHC-Fr, ICC/IF, Rat)

Chen K, Liu M, Liu Y et al. Formylpeptide receptor-2 contributes to colonic epithelial homeostasis, inflammation, and tumorigenesis J Clin Invest 2013 Mar 1 [PMID: 23454745] (ICC/IF, Mouse)

Singh D, Qi R, Jordan JL et al. The Human Antimicrobial Peptide LL-37, but Not the Mouse Ortholog, mCRAMP, Can Stimulate Signaling by Poly(I:C) through a FPRL1-dependent Pathway. J Biol Chem 2013 Mar 22 [PMID: 23386607] (WB, Human, Mouse)

## Procedures

### Immunohistochemistry-Paraffin protocol for FPRL1 Antibody (NLS1878)

#### Immunohistochemistry-Paraffin Embedded Sections

##### Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

##### Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

### Western Blot protocol for FPRL1 Antibody (NLS1878)

#### Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
  2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
  3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
  4. Rinse the blot.
  5. Block the membrane using standard blocking buffer for at least 1 hour.
  6. Wash the membrane in wash buffer three times for 10 minutes each.
  7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
  8. Wash the membrane in wash buffer three times for 10 minutes each.
  9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
  10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
  11. Apply the detection reagent of choice in accordance with the manufacturers instructions.
- Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



**Immunocytochemistry/Immunofluorescence Protocol for FPRL1 Antibody (NLS1878)**

## Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





### **Novus Biologicals USA**

8100 Southpark Way, A-8  
Littleton, CO 80120  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
novus@novusbio.com

### **Novus Biologicals Canada**

461 North Service Road West, Unit B37  
Oakville, ON L6M 2V5  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada@novusbio.com

### **Novus Biologicals Europe**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: technical@novusbio.com  
Orders: orders@novusbio.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

**For more information on our guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee).**

