


ONPG β -Galactosidase Assay Kit

INSTRUCTION MANUAL



OZBIOSCIENCES
The art of delivery systems

www.ozbiosciences.com

ONPG β -Galactosidase Assay Kit

Instruction Manual

Ready-to-use assay system for quantitatively measuring β -galactosidase expression levels in transfected cells using ONPG as a sensitive substrate.

Catalog Number: GO - 10001

You can order this product by contacting us. For all other supplementary information, do not hesitate to contact our dedicated technical support (tech@ozbiosciences.com).

OZ Biosciences SAS

163 avenue de Luminy
Case 922, zone entreprise
13288 Marseille cedex 09 - FRANCE
Ph: +33 (0) 486 948 516
Fax: +33 (0) 486 948 515
contact@ozbiosciences.com
order@ozbiosciences.com

OZ Biosciences INC

4901 Morena Blvd,
Suite 501
San Diego CA 92117 - USA
Ph : + 1-858-246-7840
Fax : + 1-855-631-0626
contactUSA@ozbiosciences.com
orderUSA@ozbiosciences.com

www.ozbiosciences.com

1. Technology

1.1. Description

Congratulations on your purchase of the ONPG β -galactosidase assay kit!

LacZ is one of the most frequently used reporter gene in transfection experiments because the gene product specific properties. Indeed, the LacZ encoded protein, β -galactosidase, is very stable, resistant to proteolytic degradation and easily tested. The levels of active β -galactosidase expression can be quickly measured by its catalytic hydrolysis of *o*-nitrophenyl- β -D-galactopyranoside (ONPG) substrate to a bright yellow product. All the necessary reagents provided in this assay kit offer a rapid, simple and sensitive method to quantify the enzyme expression level in transfected cells.

This ONPG β -galactosidase assay kit is:

- Simple and Rapid
- Ready-to-use
- Economical

1.2. Kit Contents

The kit is provided with sufficient reagents to perform 500 micro assays in 96-well plate.

Component	Quantity	Storage
5X Lysis Buffer	55 ml	4°C
Standard Dilution Buffer	55 ml	4°C
ONPG Substrate Solution (<i>o</i> -nitrophenyl- β -D-galactopyranoside)	55 ml	4°C
Stop Buffer	55 ml	4°C
β -gal enzyme standard, 40 units	100 μ L	-20°C

Stability and Storage

Storage Upon receipt and for long-term use, store all reagent tubes at the indicated storage conditions (see table above). Kit's components are stable for at least 1 year at the recommended storage temperature.

Shipping condition: The ONPG β -galactosidase assay kit is shipped with gel pack (4°C)

2. Applications and Protocols

2.1. Usage

1. Transfect cells with a plasmid expressing Lac Z gene
2. Lyse the cells using the lysis buffer
3. Transfer the lysate to a fresh tube or a micro titer plate. Dilute the lysate if needed
4. Prepare a β -galactosidase standard curve with standard dilution buffer
5. Add the substrate and monitor the color development at 405 nm - 420 nm
6. Calculate the expression levels based on a standard curve

2.2. General Conditions

- Before use, dilute the 5X Lysis buffer to 1X with distilled water. The surplus of unused 1X Lysis Buffer may be stored at +4°C for future use.
- The ONPG substrate solution is ready-to-use. **CAUTION:** Wear Gloves for manipulating the ONPG since it will stain exposed skin.

2.3. General Protocol

- **Harvesting adherent cells:**
 1. Aspirate the growth medium 24-72 hours after transfection from the culture dish including the control cells (non-transfected). Cells can be optionally washed once with 1X PBS.
 2. Add 1X Lysis Buffer to the culture dish. Solution volumes recommended for various plates are:

Type of culture dish	Volume of 1X Lysis Buffer (μ L / well)
96-well plate	50
24-well plate	250
12-well plate	500
6-well plate	1000
60 mm dish	2500
100 mm dish	5000

3. Incubate the dish 10-15 min. at room temperature by swirling it slowly several times to ensure complete lysis. The dishes can be observed under a microscope to confirm the complete lysis.

NOTE: A fast freeze/thaw cycle (freeze 1-2 hours at -20°C or -70°C and thaw at room temperature) of the dish can also be done to achieve a good lysis. Proceed to the colorimetric assay or freeze the plate at -70°C until ready.

4. **OPTIONAL:** The dish can be centrifuged for 2-3 minutes to pellet the insoluble material before proceeding to the colorimetric assay. Then, the supernatant is ready to be tested.

- **Harvesting suspension cells:**

1. 24-72 hours post-transfection, centrifuge the cells at 250 x g for 5 minutes. Then, aspirate the supernatant. Cells pellet can be optionally washed once with 1X PBS.
2. Resuspend the cell pellet in 1X Lysis Buffer. The amount of Lysis Buffer depends on the size of the culture dishes used for transfection (i.e., cell pellet size) and we recommend using between 50 to 2000 L.
3. Incubate the cell lysate 10-15 min. at room temperature by gently swirling the dishes several times to ensure complete lysis. Proceed to the colorimetric assay or freeze the plate at -70°C until ready.
NOTE: A quick freeze/thaw cycle (freeze 1-2 hours at -20°C or -70°C and thaw at room temperature) can also be done to obtain a good lysis.
4. **OPTIONAL:** The dish can be centrifuged 2 min. to pellet the insoluble material before proceeding to the colorimetric assay. Then, the supernatant is ready to be assayed.

2.4. 96 well micro titer plate assay*

1. If needed thaw the dish, tube or plate of lysed cells at room temperature. If the transfection is performed with a 96-well plate, perform the assay directly on the plate.
2. Add 50 µl of Standard Dilution Buffer to the wells of a 96-well plate (flat bottom) except the control wells, which are save for the standard curve.
3. Prepare a serial dilution of β-galactosidase standard in Standard Dilution Buffer. Then, transfer 50 µL of each point of the standard curve to the control wells (containing 50µL of cell lysate). A dilution protocol for an optimized standard curve is suggested in the following table :

β-Gal Standard (milliunits)	Standard Dilution Buffer Volume	β-Gal Standard Volume
200	990 µL	10 µL of β-gal standard stock
100	200 µL	200 µL of 200 mu β-gal standard
50	200 µL	200 µL of 100 mu β-gal standard
25	200 µL	200 µL of 50 mu β-gal standard
12.5	200 µL	200 µL of 25 mu β-gal standard
6.25	200 µL	200 µL of 12.5 mu β-gal standard
3.125	200 µL	200 µL of 6.25 mu β-gal standard
1.562	200 µL	200 µL of 3.125 mu β-gal standard

NOTE 1: Adjust the standard curve to suit the specific experimental conditions, such as cell type, cell number, transfection reagent, size of the culture dish or plasmid vector.

NOTE 2: The dilutions for the standard curve must be prepared freshly each time the assay is performed.

1. Add 50 µL of each sample/well.

NOTE: It may be necessary to dilute the lysate in 1X Lysis Buffer when transfection efficiency is very high. In contrast, when transfection efficiency is low, reduce the volume of lysis buffer used to harvest the cells

(see above) or use up to 150 μL of cell lysate for the colorimetric assay. If the transfection is performed with a 96-well plate, perform the assay directly on the plate.

To control endogenous β -galactosidase activity, prepare controls (blank) by adding 50 μL of lysis buffer to a well and 50 μL of cell lysate from non-transfected cells to another well.

2. Add 100 μL of ONPG Substrate Solution to each well. Incubate the plate at room temperature until the yellow color develops (from approximately 10 min. to 4 h depending on the cell type).
3. Read the absorbance at 405-420 nm with a micro titer spectrophotometer. Stop solution is not required for this format, since all wells are read simultaneously. Avoid the presence of bubbles in the wells while reading. Bubbles will interfere with the absorbance reading and can be removed with a fine gauge needle, tips or very weak gas flow.
4. Quantify β -galactosidase expression based on a linear standard curve.

*Felgner, J.H. *et al.* Enhanced gene delivery and mechanism studies with a novel series of cationic lipid formulations. *J. Biol. Chem.* **269**, 2550-2561 (1994)

2.5. Macro assay

1. Thaw the cell lysate (if needed) and transfer 100 μL to a fresh tube, or 50 μL to a 96-well plate. If a 96-well plate is used, follow the protocol described above.

NOTE: It may be necessary to dilute the cell lysate in 1X Lysis Buffer when transfection efficiency is very high. In contrast, when transfection efficiency is very low, reduce the volume of lysis buffer used to harvest the cells (see description above) or use up to 150 μL of cell extract for the colorimetric assay

Prepare a blank by adding 100 μL of lysis buffer to a tube. Add also 100 μL of cell lysate from non-transfected cells (mock-transfected cells) to a tube to control endogenous β -galactosidase activity.

2. Add 50 μL of Standard Dilution Buffer to each tube.
3. Prepare a serial dilution of β -galactosidase (E. Coli) standards with Standard Dilution Buffer separately. Transfer 50 μL of each standard to a fresh tube containing 100 μL cell lysate from non-transfected cells. The highest recommended amount of beta-galactosidase is 400,000 pg. (200 milliunits). Adjust the standard curve to suit the specific experimental conditions, such as cell type, transfection reagent, or plasmid vector. 2X serial dilution of standard curve consisting of 8 points is recommended. A dilution protocol example is shown in the section of 96-well plate assay.
4. Add 300 μL of the ONPG Substrate Solution to each tube. Incubate the tubes at room temperature until the yellow color develops (from approximately 10 minutes to 4 hours depending on the cell type). Add 500 μL of Stop Solution to stop the reaction. Final volume is 950 μL .
5. Read the absorbance at 405-420 nm with a spectrophotometer.
6. Quantify β -galactosidase expression based on a linear standard curve.

3. Related Products

Description
MAGNETOFECTION TECHNOLOGY
Super Magnetic Plate (<i>standard size for all cell culture support</i>)
Mega Magnetic plate (<i>mega size to hold 4 culture dishes at one time</i>)
Transfection reagents:
PolyMag Neo (<i>for all nucleic acids</i>)
SilenceMag (<i>for siRNA application</i>)
NeuroMag (<i>dedicated for neurons</i>)
Transfection enhancer:
CombiMag (<i>to improve any transfection reagent efficiency</i>)
Viral Transduction enhancers:
ViroMag (<i>to optimize viral transduction</i>)
ViroMag R/L (<i>specific for retrovirus and Lentivirus</i>)
AdenoMag (<i>for Adeno viruses</i>)
LIPOFECTION TECHNOLOGY (LIPID-BASED)
Lullaby (<i>siRNA transfection reagent</i>)
DreamFect Gold (<i>Transfection reagent for all types of nucleic acids</i>)
Ecotransfect (<i>Economical reagent for routine transfection</i>)
FlyFectin (<i>for Insect cells</i>)
VeroFect (<i>for Vero cells</i>)
3D TRANSFECTION TECHNOLOGY
3Dfect (<i>for scaffolds culture</i>)
3DfectIN (<i>for hydrogels culture</i>)
RECOMBINANT PROTEIN PRODUCTION
HYPE-5 Transfection Kit (<i>for High Yield Protein Expression</i>)
PROTEIN DELIVERY SYSTEMS
Ab-DeliverIN (<i>delivery reagent for antibodies</i>)
Pro-DeliverIN (<i>delivery reagent for protein in vivo and in vitro</i>)
PLASMIDS PVECTOZ
pVectOZ-LacZ 25µg
pVectOZ-SEAP 25µg
ASSAY KITS
Bradford – Protein Assay Kit
β-Galactosidase assay kits (CPRG/ONPG)
X-Gal Staining Kit
BIOCHEMICALS
D-Luciferin, K ⁺ and Na ⁺ 1g
G-418, Sulfate 1g
X-Gal powder 1g

Do not hesitate to contact us for all complementary information and remember to visit our website in order to stay informed on our last breakthrough technologies and updated on our complete product list.

Purchaser Notification

Limited License

The purchase price paid for the ONPG β -galactosidase assay kit by end users grants them a non-transferable, non-exclusive license to use the kit and/or its separate and included components (as listed in the section 1, Kit Contents). This reagent is intended **for internal research only** by the buyer. Such use is limited to the use described in the product manual. Furthermore, research only use means that this kit and all of its contents are excluded, without limitation, from resale, repackaging, or use for the making or selling of any commercial product or service without the written approval of OZ Biosciences.

Purchasers may terminate this License at any time by returning all ONPG β -galactosidase assay kit material and documentation to OZ Biosciences, or by destroying all ONPG β -galactosidase assay kit components. Purchasers are advised to contact OZ Biosciences with the notification that a ONPG β -galactosidase assay kit is being returned in order to obtain a refund and/or to expressly terminate a research only license granted through the purchase of the kit(s).

This document covers in full the terms of the ONPG β -galactosidase assay kit research only license, and does not grant any other express or implied license. The laws of the French Government shall govern the interpretation and enforcement of the terms of this License.

Product Use Limitations

The ONPG β -galactosidase assay kit and all of its components are developed, designed, intended, and sold for research use only. They are not to be used for human diagnostic or included/used in any drug intended for human use. All care and attention should be exercised in the handling of the kit components by following appropriate research lab practices.

For more information, or for any comments on the terms and conditions of this License, please contact:

Director of Business Development
OZ Biosciences SAS
Parc Scientifique et Technologique de Luminy
Bât. Grand luminy, zone entreprise
Case 922, 13288 Marseille Cedex 9 - France
Ph: +33 (0)4.86.94.85.16
Fax: +33 (0)4.86.94.85.15
E-mail: contact@ozbiosciences.com

CONTACTS

OZ Biosciences SAS
163 avenue de Luminy
Case 922, zone entreprise
13288 Marseille cedex 09
FRANCE

Ph: +33 (0) 486 948 516
Fax: +33 (0) 486 948 515

contact@ozbiosciences.com
order@ozbiosciences.com
tech@ozbiosciences.com

OZ Biosciences INC
4901 Morena Blvd,
Suite 501
San Diego CA 92117
USA

Ph : + 1-858-246-7840
Fax : + 1-855-631-0626

contactUSA@ozbiosciences.com
orderUSA@ozbiosciences.com
techUSA@ozbiosciences.com

www.ozbiosciences.com

Follow us!

