

Instruction Manual

Product	Size	Catalog Number
Human CD14 ⁺ Monocytes from Peripheral Blood (hMoCD14 ⁺ -PB), single donor	10 x 10 ⁶ cryopreserved cells	C-12909
Human CD14 ⁺ Monocytes from Peripheral Blood (hMoCD14 ⁺ -PB), single donor, pre-screened	10 x 10 ⁶ cryopreserved cells	C-12912
Human Monocytes from Peripheral Blood (hMo-PB), single donor	10 x 10 ⁶ cryopreserved cells	C-12910
Human Monocytes from Peripheral Blood (hMo-PB), single donor, pre-screened	10 x 10 ⁶ cryopreserved cells	C-12913

Product Description

Monocytes (Mo) are immature phagocytic cells circulating in the blood. Acting as antigen-presenting immune cells, they can phagocytize and degrade microbes and particulate matter. Monocytes can differentiate into several different cell types. The most common applications for Monocytes are the *in vitro* differentiation into monocyte-derived dendritic cells (moDC), macrophages, and osteoclasts as well as their use as phagocytic immune cells in immunology and infection biology.

CD14 belongs to the family of LPS receptor antigens and is strongly expressed on the majority of Monocytes. Thus, binding of immunomagnetic particles to this antigen is employed for purification of PromoCell CD14⁺-Monocytes (hMoCD14⁺-PB) from Mononuclear Cells (MNC).

In contrast, the “untouched” Monocytes (hMo-PB) provided by PromoCell are enriched from MNC by a density gradient-based method completely devoid of using any immunomagnetic beads. The advantage of these completely native Monocytes is that poten-

tial effects of bound magnetic particles (which can be ingested by this phagocytic cell type) are excluded.

In addition, all PromoCell Monocytes are available as pre-screened cells, which have been positively tested for their differentiation capacity into dendritic cells. PromoCell Human CD14⁺ Monocytes and PromoCell Human “untouched” Monocytes are isolated from fresh peripheral blood. After isolation of the ultra-pure mononuclear cell fraction by proprietary methods, the CD14⁺ Monocytes are isolated by positive selection and the “untouched” Monocytes are purified by an advanced multi-gradient technique.

Immediately after isolation, the freshly prepared human monocytes are cryopreserved using PromoCell’s proprietary serum-free, freezing medium Cryo-SFM. Each cryo vial contains more than 10 million viable cells after thawing.

Quality Control

Rigid quality control tests are performed for each lot of PromoCell Monocytes. They are routinely characterized by flow cytometry analyzing the viability, cell

size and purity.

In addition, all cells have been tested for the absence of HIV-1, HIV-2, HBV and HCV, and microbial contaminants (fungi, bacteria, and mycoplasma).

A detailed certificate of analysis (CoA) for each lot can be downloaded at: www.promocell.com/coa

Intended Use

PromoCell Monocytes are for *in vitro* research use only and not for diagnostic or therapeutic procedures.

Warning

Although tested negative for HIV-1, HIV-2, HBV, and HCV, the cells - like all products of human origin - should be handled as potentially infectious. No test procedure can completely guarantee the absence of infectious agents.

Follow appropriate safety precautions!

After delivery, start immediately with the protocol for cryopreserved cells (see page 2).

*Start immediately after delivery.
Use aseptic techniques and a laminar flow bench.*

Protocol for Cryopreserved Cells

Straight after arrival, store the cryopreserved cells in liquid nitrogen, or seed them immediately.

Note: Storage at -80°C is not sufficient for cell preservation and causes irreversible cell damage.

1. Prepare the medium

Refer to the recommended seeding density (see page 4) and the cell number per vial given in the lot-specific CoA of the cells. Transfer the needed volume of PromoCell Medium in cell culture vessels. For equilibration, place the vessels in an incubator (37°C , 5% CO_2) for 30 minutes.



2. Thaw the cells

Remove the cryovial from the liquid nitrogen container and immediately place it on dry ice - even for short transportation. Under a laminar flow bench, briefly twist the cap a quarter turn to relieve pressure, then re-tighten. Immerse the vial into a water bath (37°C) just up to the screw cap for 2 minutes. Ensure that no water enters the thread of the screw cap.



3. Disinfect the vial and seed the cells

Thoroughly rinse the cryovial with 70% ethanol under a laminar flow bench. Then, aspirate the excess ethanol from the thread area of the screw cap. Open the vial and transfer the cells with a 2 ml serological pipette (not a micropipette) to a cell culture vessel containing the pre-warmed medium from step 1 without resuspending.

Note: Do not resuspend the cells at any time, since clumping may occur.



4. Incubate the cells

Place the vessel in an incubator (37°C , 5% CO_2). For complete recovery, leave the cells untouched for at least 18 hours. Do not disrupt the flask during this recovery stage. Change the medium after 18 - 24 hours.

Note: Handling of the cells before complete recovery results in clumping.




Use aseptic techniques and a laminar flow bench.

Subcultivation Protocol


1. Harvest the cells

Harvest the cell suspension and determine the cell number. Spin down the cells for 10 minutes at 240 x g.



2. Resuspend and reseed cells

Discard the supernatant (step 1), add 1 ml of the appropriate PromoCell Medium (step 2), and resuspend the cells by carefully pipetting up and down. Seed the cells according to the recommended seeding density in new cell culture vessels containing fresh PromoCell Medium prewarmed to 37°C. Place the vessels in an incubator (37°C, 5% CO₂).



Specifications

Product	Recommended Culture Media	Plating density
Human CD14 ⁺ Monocytes from Peripheral Blood (hMoCD14 ⁺ -PB), single donor	C-28030	1 x 10 ⁶ cells per ml
Human CD14 ⁺ Monocytes from Peripheral Blood (hMoCD14 ⁺ -PB), single donor, pre-screened	C-28030 C-28050	1 x 10 ⁶ cells per ml
Human Monocytes from Peripheral Blood (hMo-PB), single donor	C-28030	1 x 10 ⁶ cells per ml
Human Monocytes from Peripheral Blood (hMo-PB), single donor, pre-screened	C-28030 C-28050	1 x 10 ⁶ cells per ml

Related Products

Product	Size	Catalog Number
Mononuclear Cell Medium (Ready-to-use)	500 ml	C-28030
DC Generation Medium	250 ml	C-28050
DC Base Medium	250 ml	C-28053
Cryo-SFM	30 ml	C-29910
	125 ml	C-29912
hMNC-PB pooled Pellet	1 million cells per pellet	C-14099
hMoCD14 ⁺ -PB single donor Pellet	1 million cells per pellet	C-14110
hMoCD14 ⁺ -PB single donor, pre-screened Pellet	1 million cells per pellet	C-14112
hMo-PB single donor Pellet	1 million cells per pellet	C-14111
hMo-PB single donor pre-screened Pellet	1 million cells per pellet	C-14113
GM-CSF, human, recombinant	10 µg	C-60420
IL-4 CC, human, recombinant	5 µg	C-61401
M-CSF, human, recombinant	10 µg	C-60442
sRANKL, human, recombinant	10 µg	C-63200

PromoCell GmbH

Sickingenstr. 63/65
69126 Heidelberg
Germany

Email: info@promocell.com
www.promocell.com

USA/Canada

Phone: 1 – 866 – 251 – 2860 (toll free)
Fax: 1 – 866 – 827 – 9219 (toll free)

Deutschland

Telefon: 0800 – 776 66 23 (gebührenfrei)
Fax: 0800 – 100 83 06 (gebührenfrei)

France

Téléphone: 0800 90 93 32 (ligne verte)
Téléfax: 0800 90 27 36 (ligne verte)

United Kingdom

Phone: 0800 – 96 03 33 (toll free)
Fax: 0800 – 169 85 54 (toll free)

Other Countries

Phone: +49 6221 – 649 34 0
Fax: +49 6221 – 649 34 40