


5,5',6,6'-tetrachloro-1,1',3,3'-tetraethyl-benzimidazolylcarbocyanine iodide

Instruction Manual

Catalog Number	PK-CA707-70014
Description	JC-1 is a mitochondrial dye that stains mitochondria in living cells in a membrane potential-dependent fashion. JC-1 monomer is in equilibrium with so-called J-aggregates, which are favored at higher dye concentration or higher mitochondrial membrane potential. The monomer JC-1 has green fluorescence ($\lambda_{em} = 527 \text{ nm}$), while the J-aggregates have red fluorescence ($\lambda_{em} = 590 \text{ nm}$). Therefore, it has been possible to use fluorescence ratioing technique to study mitochondrial membrane potentials. JC-1 is particularly useful for apoptosis studies. In apoptotic cells, the dye stays in the cytoplasm and fluoresces green. It has also been applied in high throughput drug screening applications. Extinction coefficient (in MeOH, 505 nm) = 190,000. PromoKine offers JC-1 in both chloride salt (PK-CA707-70011) and iodide salt (PK-CA707-70014) forms. Both forms have identical spectral properties. The iodide form has been used in most of the publication. However, some researchers may prefer the chloride form since Cl ⁻ is the most prevalent anion in biological systems.
Quantity	5 mg
Excitation / Emission Maxima	see "Description"
Molecular Structure	
Molecular Weight / Molecular Formula	652 Da; C ₂₅ H ₂₇ Cl ₄ IN ₄
Purity	>99% (as determined by HPLC)
Appearance / Formulation / Solubility	Red solid; soluble in DMSO.
Storage & Stability	Store desiccated at 4°C. Protect from light, especially when in solution.
Applications	See Description
References	<p>1) Smiley, S.T., et al. Proc. Natl. Acad. Sci. 88, 3671(1991)</p> <p>2) Reers, M., et al. Biochemistry 30, 4480(1991).</p> <p>References on use for apoptosis studies:</p> <p>1) "Use of Flow Cytometry techniques in Studying Mechanisms of Apoptosis in Leukemic Cells", A.M. Gorman, et al. Cytometry 29, 97(1977)</p> <p>2) "JC-1, but Not DiOC6(3) or Rhodamine 123, is a Reliable Fluorescent Probe to Assess Changes in Intact Cells: Implications for Studies on Mitochondrial Functionality during Apoptosis." S. Salvioli, et al. FEBS Lett. 411, 77(1997)</p> <p>3) "Functional Assay of Multidrug Resistant Cells Using JC-1, a Carbocyanine Fluorescent Probe." J.M. Kuhnle, et al. Leukemia, 11, 1147(1997).</p>
Caution	Potentially harmful. Avoid prolonged or repeated exposure. Avoid getting in eyes, on skin, or on clothing. Wash thoroughly after handling. If eye or skin contact occurs, wash affected areas with plenty of water for 15 minutes and seek medical advice. In case of inhaling or swallowing, move individual to fresh air and seek medical advice immediately.

FOR IN VITRO RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC PROCEDURES.