

# Augurin, Prepro (71-107) (Human) - Antibody for Immunohistochemistry

Catalog#	Standard size	Price	Order
H-012-22	100 µl	\$250.00	<input type="button" value="ORDER"/>


## Product description

## More information


[Augurin - Research Abstracts, Antibody Stainings, Kits Standard Curves, Sequences, etc.](#)

[Isolation and Identification of Augurin Peptide from the Bovine Hypothalamus](#)

## Image



Human pancreas tissue was stained by rabbit anti-Augurin (71-107) (H) Antiserum (Cat. No.: H-012-22)



Human pancreas tissue was stained by rabbit anti-Augurin, Prepro (71-107) (H) Antiserum (Cat. No.: H-012-22)

## Protocol for Immunohistochemistry

Tissue Sample	Human pancreas tissues
Fixative	10% Formalin
Embedding	Paraffin
Negative control	No primary antibody
Pretreatment	N/A
Blocking	3% H <sub>2</sub> O <sub>2</sub> , 2% Normal Goat Serum
Primary Antibody	Rabbit Anti-Augurin, Prepro (71-107) (H) Antiserum (Catalog No.: H-012-22)
Optimal Dilution	1:500, 1 hour at RT
Secondary Antibody	Goat anti-Rabbit IgG, Biotinylated (1:400, 30 min)
Amplification	Streptavidin-HRP (Vector), 1:400, 30 min
Detection system	HRP
Substrate	DAB (Sigma), 3 min
Counterstained	Hematoxylin, 30 Sec

**Storage Condition** For optimal results, use the antibody as soon as possible after reconstitution. Store in lyophilized form unless needed and reconstitute immediately before use. Once reconstituted, the antibody should be stable for a few days at -4°C. For storage up to a few months, prepare small aliquots after reconstitution and freeze at -20°C or -80°C. Repeated freeze thaw cycles should be strictly avoided.

**Recommended Dilution Factor** Immunohistochemistry - 1:500

Thursday 20 December, 2012

11986430 requests since Wednesday 05 April, 2006

Copyright © 2012 PHOENIX PHARMACEUTICALS, INC.