GLP-2 (1-34) (Human) - Antibody for Immunohistochemistry

Catalog# Standard size Order ORDER H-028-14 50 µl \$150.00 Product description More information References


GLP-1 & GLP-2

Normal Plasma Peptide/Protein Levels

Neuron Function Modulation

I mage





Protocol for Immunohistochemistry

Tissue Sample **Human Brain Tissue**

Fixative 10% Formalin

Paraffin **Embedding**

Negative control No primary antibody

Pretreatment Target retrieval 25 min (Steam) Blocking 3% H₂O₂, 2% Normal Goat Serum

Primary Antibody Rabbit Anti-GLP-2 (H) Antiserum (Catalog No.: H-028-14)

Optimal Dilution 1:250~500, 1 hour at RT

Secondary Antibody Goat anti-Rabbit IgG, Biotinylated (1:400, 30 min)

Amplification Streptavidin-HRP (Vector), 1:400, 30 min

Detection system HRP

Substrate DAB (Sigma), 3 min Counterstained Hematoxylin, 30 Sec

Catalog # H-028-14

Standard Size 50 µl

Sequence His - Ala - Asp - Gly - Ser - Phe - Ser - Asp - Glu - Met - Asn - Thr - Ile - Leu - Asp - Asn - Leu - Ala - Ala -

Arg - Asp - Phe - Ile - Asn - Trp - Leu - Ile - Gln - Thr - Lys - Ile - Thr - Asp - Arg

Species Human

Host Rabbit

For best and reproducible results, reconstitute with 50µl of distilled water for the equivalent of undiluted Reconstitution

antiserum, immediately before use. Do not refreeze any unused portions.

Storage Condition

For optimal results, use the antibody as soon as possible after reconstitution. Store in lyophilized form unless needed and reconstitute immediately before use. Once reconstituted, the antibody should be stable for a few days at -4°C. For storage up to a few months, prepare small aliquots after reconstitution and

freeze at -20°C or -80°C. Repeated freeze thaw cycles should be strictly avoided.

Content This vial contains 50µl of Rabbit Anti- Glucagon-Like Peptide-2 (GLP-2) (Human) Serum in the lyophilized

form.