

CCK (46-69) (Human) - Antibody for Immunohistochemistry

Catalog#	Standard size	Price	Order
H-069-27	100 ul	\$350.00	<input type="button" value="ORDER"/>


Product description

More information

[All About Cholecystokinin](#)

[Normal Plasma Peptide/Protein Levels](#)

Image



Human colon cancer tissue was stained by Rabbit Anti-CCK (46-69) (H) Antibody (Cat. No.: H-069-27)

Protocol for Immunohistochemistry

Tissue Sample	Human colon cancer tissues
Fixative	10% Formalin
Embedding	Paraffin
Negative control	No primary antibody
Pretreatment	N/A
Blocking	3% H ₂ O ₂ , 2% Normal Goat Serum
Primary Antibody	Rabbit anti-CCK (46-69) (Human) antibody (Cat. No.: H-069-27)
Optimal Dilution	1:500
Secondary Antibody	Goat Anti-Rabbit IgG, Biotinylated (1:400), 30 min
Amplification	Streptavidin-HRP (Vector), 1:400, 30 min
Detection system	HRP
Substrate	DAB (Sigma), 3 min
Counterstained	Hematoxylin, 30 Sec

Catalog #	H-069-27
Family	CCK
Standard Size	100 ul
Sequence	Val - Ser - Gln - Arg - Thr - Asp - Gly - Glu - Ser - Arg - Ala - His - Leu - Gly - Ala - Leu - Leu - Ala - Arg - Tyr - Ile - Gln - Gln - Ala
Species	Human
Host	Rabbit
Immunogen	CCK (46-69) (Human)
Reconstitution	For best and reproducible results, reconstitute with 100µl of distilled water for the equivalent of undiluted antiserum, immediately before use. Do not refreeze any unused portions.
Storage Condition	For optimal results, use the antibody as soon as possible after reconstitution. Store in lyophilized form unless needed and reconstitute immediately before use. Once reconstituted, the antibody should be stable for a few days at -4°C. For storage up to a few months, prepare small aliquots after reconstitution and freeze at -20°C or -80°C. Repeated freeze thaw cycles should be strictly avoided.