
Human RXR alpha stable cells (Inducible)

Species:	<i>Homo Sapiens</i>	Catalog Number:	CLHKRXR02
Cell Number:	~2x10 ⁶	Storage/Shipping:	Dry Ice/-80°C

Introduction:

Nuclear hormone receptors (NHRs) are a superfamily of transcription factors that function as powerful metabolic regulators to control a variety of systemic processes in physiology. They also play key roles in the pathophysiology of many major disease states, such as diabetes, obesity, inflammation, atherosclerosis and heart failure. To date, this superfamily has provided a rich source of drug design targets and it is continuing to be one of the hottest areas for pharmaceutical researches.

PrimCells now provides a comprehensive set of NHR expressing cell lines to the research community. These high-quality, flag-tagged NHR-expressing cell lines aim to facilitate further biochemical and molecular studies of their functions and hence encourage new strategies for drug design.

The human RXR alpha stable cell line was created by stable co-transfection of HEK293 cells with a plasmid constitutively expressing HA tagged the human RXR alpha protein.

Thawing of Frozen Cells

1. Upon receipt of the frozen cells, it is recommended to thaw the cells and initiate the culture immediately in order to retain the highest cell viability.
2. To thaw the cells, put the vial in 37°C water bath with gentle agitation for ~1min. Keep the cap out of water to minimize the risk of contamination.
3. Pipette the cells into a 15ml conical tube with ~5ml fresh culture medium.
4. Centrifuge at 1000rpm (~220g) for 5min under room temp.
5. Remove the supernatant and resuspend the cells in fresh culture medium
6. Transfer the cells into new tissue culture flasks and move them to 37°C incubator (5% CO₂) for continuous culture.

Safety Precaution: *it is highly recommended that protective gloves and clothing should be used when handling frozen vials.*

Standard Culture Procedure

1. Cells should be maintained in the complete culture medium until reaching ~80-90% confluence. **Note:** *Never let the cells to become over confluent.*
2. Add ~2.5ml of 0.05% Trypsin-EDTA to the flask and incubate for 5min at 37°C.

3. Neutralize the enzyme activity by adding 2-3 volumes of fresh complete culture medium.
4. Centrifuge 1000rpm (~220g) for 5min and resuspend the cells in desired volume of medium.
5. Transfer the cells to a new tissue culture treated flask for subculture. **Note:** *It is recommended that cells are passaged at the ratio of 1:10.*

Complete Growth Medium

DMEM (Corning, Cat#10014CV): 450ml

FBS: 50ml

Total Volume: 500ml

Technical Support

For additional information regarding the product and technical questions, please contact Supports@PrimCells.com. You are guaranteed to receive a response within 24hrs from one of our scientists.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While PrimCells uses reasonable efforts to include accurate and up-to-date information on this product sheet, PrimCells makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. PrimCells does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. PrimCells is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, PrimCells is not liable for damages arising from the misidentification or misrepresentation of cultures.

© PrimCells 2013. All rights reserved.