


Sulfo-NHS-Biotin Protocol and Product Information Sheet

Product Category:	Biotinylation Reagents
Catalog Number(s):	b2100-100mg , b2100-1gm , b2100-custom
Product Name:	Sulfo-NHS-Biotin
Alternative Name(s):	Sulfosuccinimidyl biotin; Sulfosuccinimido biotin; Biotin 3-sulfo-N-hydroxysuccinimide ester sodium salt; N-hydroxysulfosuccinimidyl biotin
CAS Number:	119616-38-5
Chemical Formula:	C ₁₄ H ₁₈ N ₃ NaO ₈ S ₂
Molecular Weight:	443.43
Spacer Length:	13.5 Å



General Sulfo-NHS-Biotin Protein Biotinylation Protocol

1. Allow vial of Sulfo-NHS-Biotin to equilibrate to ambient temperature before opening.
2. Dissolve protein at a concentration of 1-10 mg/mL in 100 mM sodium phosphate, 150 mM NaCl, pH 7.2-7.5 or other suitable amine-free buffer.
3. Immediately before use, create a 20 mg/mL Sulfo-NHS-Biotin stock solution in water or buffer (same as step 2). Anhydrous DMF ([cr8106-25ml](#)) or DMSO ([cr8105-25ml](#)) can be used to make a stock solution ahead of time.
4. Add sufficient Sulfo-NHS-Biotin stock solution to the protein solution to obtain 10-20 fold molar excess of biotinylation reagent over protein.
Note: Alternatively, an amount of Sulfo-NHS-Biotin can be added to the protein solution required to give 10-20 fold molar excess. Dilute protein solutions (i.e. 1-2 mg/mL) may require increased molar excess of Sulfo-NHS-Biotin (i.e. ≥ 20 fold) to yield similar biotinylation of a more concentrated protein solution.
5. Allow biotinylation reaction to proceed for 30-60 minutes at room temperature or ≥ 2 hours at 4°C.
6. Desalt biotinylated protein through dialysis or gel filtration with a resin, such as Sephadex® G-25 ([q4109](#)) or equivalent.

References:

Hermanson, G.T. 1996. Bioconjugate Techniques. Academic Press, San Diego, CA, USA.