



# **Certificate of Analysis**

www.tocris.com

Product Name: (S)-(-)-Carbidopa Catalog No.: 0455 Batch No.: 4

CAS Number: 28860-95-9 EC Number: 249-271-9

IUPAC Name: (S)- $\alpha$ -Hydrazino- $\alpha$ -methyl- $\beta$ -(3,4-dihydroxyphenyl)propionic acid

### 1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula:  $C_{10}H_{14}N_2O_4.H_2O$ 

Batch Molecular Weight: 244.25
Physical Appearance: White solid
Solubility: DMSO to 20 mM

Solubility: DMSO to 20 ml Storage: Store at RT

Batch Molecular Structure:

HO Me NHNH<sub>2</sub>

2. ANALYTICAL DATA

HPLC: Shows 99.4% purity

1H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

**Optical Rotation:**  $[\alpha]_D = -18.5$  (Concentration = 1, Solvent = Methanol)

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 49.18 6.6 11.47 Found 48.96 6.52 11.5





## **Product Information**

www.tocris.com

Product Name: (S)-(-)-Carbidopa Catalog No.: 0455 Batch No.: 4

CAS Number: 28860-95-9 EC Number: 249-271-9

IUPAC Name: (S)- $\alpha$ -Hydrazino- $\alpha$ -methyl- $\beta$ -(3,4-dihydroxyphenyl)propionic acid

#### **Description:**

Peripheral decarboxylase inhibitor, used in combination with levodopa for treatment of Parkinsonism.

## **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>10</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>.H<sub>2</sub>O

Batch Molecular Weight: 244.25 Physical Appearance: White solid

**Minimum Purity: >99%** 

#### **Batch Molecular Structure:**

Storage: Store at RT

#### Solubility & Usage Info:

DMSO to 20 mM

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

#### References:

Merck Index 12 1843.

**Kopfhammer** *et al* (1985) L-Dopa infusion experiments in combination with the L-dopa decarboxylase inhibitors benserazide and carbidopa - L-dopa metabolism and neuroendocrinological effects. Pharmacopsychiatry *18* 143.

Cheng et al (1992) Carbidopa inhibition of aromatic L-amino acid decarboxylase in rat brain capilliaries. FASEB J. 6 A1589.

