


Recombinant SARS-CoV-2 Spike RBD B.1.617.2 L452R T478K His-tag

Catalog Number: 10876-CV

DESCRIPTION	
Source	Human embryonic kidney cell, HEK293-derived sars-cov-2 Spike RBD protein Arg319-Phe541 (Leu452Arg, Thr478Lys), a C-terminal 6-His tag Accession # YP_009724390.1
N-terminal Sequence Analysis	Arg319
Predicted Molecular Mass	26 kDa

SPECIFICATIONS	
SDS-PAGE	30-37 kDa, under reducing conditions.
Activity	Measured by its binding ability in a functional ELISA with Recombinant Human ACE-2 His-tag (Catalog # 933-ZN).
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE	
Reconstitution	Reconstitute at 500 μg/mL in PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
	 12 months from date of receipt, -20 to -70 °C as supplied.
	 1 month, 2 to 8 °C under sterile conditions after reconstitution.
	 3 months, -20 to -70 °C under sterile conditions after reconstitution.



Rev. 6/17/2021 Page 1 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 **Canada** TEL 855 668 8722 **China** TEL +86 (21) 52380373 **Europe | Middle East | Africa** TEL +44 (0)1235 529449



Recombinant SARS-CoV-2 Spike RBD B.1.617.2 L452R T478K His-tag

Catalog Number: 10876-CV

BACKGROUND

SARS-CoV-2, which causes the global pandemic coronavirus disease 2019 (Covid-19), belongs to a family of viruses known as coronaviruses that also include MERS and SARS-CoV-1. Coronaviruses are commonly comprised of four structural proteins: Spike protein (S), Envelope protein (E), Membrane protein (M) and Nucleocapsid protein (N) (1). The SARS-CoV-2 S protein is a glycoprotein that mediates membrane fusion and viral entry. The S protein is homotrimeric, with each ~180-kDa monomer consisting of two subunits, S1 and S2 (2). In SARS-CoV-2, as with most coronaviruses, proteolytic cleavage of the S protein into S1 and S2 subunits is required for activation. The S1 subunit is focused on attachment of the protein to the host receptor while the S2 subunit is involved with cell fusion (3-5). A metallopeptidase, angiotensin-converting enzyme 2 (ACE-2), has been identified as a functional receptor for SARS-CoV-2 through interaction with a receptor binding domain (RBD) located at the C-terminus of S1 subunit (6, 7). Based on amino acid (aa) sequence homology, the SARS-CoV-2 S1 subunit RBD has 73% identity with the RBD of the SARS-CoV-1 S1 RBD, but only 22% homology with the MERS S1 RBD. The SARS-CoV-2 delta variant (B.1.617.2) carrying the amino acid substitution L452R and T478K in the RBD was identified as a prevalent strain in India and has been detected in more than 40 countries (8, 9). It has higher transmissible rate and more resistant to vaccine (10).

References:

- 1. Wu, F. et al. (2020) Nature 579:265.
- 2. Tortorici, M.A. and D. Veesler (2019) Adv. Virus Res. 105:93.
- 3. Bosch, B.J. et al. (2003) J. Virol. 77:8801.
- 4. Belouzard, S. et al. (2009) Proc. Natl. Acad. Sci. 106:5871.
- 5. Millet, J.K. and G.R. Whittaker (2015) Virus Res. 202:120.
- 6. Li, W. et al. (2003) Nature 426:450.
- 7. Wong, S.K. et al. (2004) J. Biol. Chem. 279:3197.
- 8. Yadav, P.D. et al. (2021) bioRxiv https://doi.org/10.1101/2021.04.23.441101.
- 9. Cherian, S. et al. (2021) bioRxiv https://doi.org/10.1101/2021.04.22.440932.
- 10. Bernal, J. et al. (2021) medRxiv https://doi.org/10.1101/2021.05.22.21257658.

Rev. 6/17/2021 Page 2 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449