



Certificate of Analysis

www.tocris.com

Product Name: Clemastine fumarate Catalog No.: 1453 Batch No.: 2

CAS Number: 14976-57-9 EC Number: 239-055-2

IUPAC Name: (2R)-2-[2-[(1R)-1-(4-Chlorophenylethoxy]ethyl]-1-methyl-2-pyrrolidine fumarate

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{21}H_{26}CINO.C_4H_4O_4$

Batch Molecular Weight: 459.97
Physical Appearance: White solid

Solubility: DMSO to 50 mM

ethanol to 10 mM

Storage: Store at RT

Batch Molecular Structure:

2. ANALYTICAL DATA

HPLC: Shows 99.9% purity

¹H NMR: Consistent with structure Mass Spectrum: Consistent with structure

Optical Rotation: $[\alpha]_D = +13$ (Concentration = 1, Solvent = Methanol)

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 65.28 6.57 3.05 Found 65.48 6.54 3.2



Product Information

Print Date: Apr 28th 2015

www.tocris.com

Product Name: Clemastine fumarate Catalog No.: 1453 Batch No.: 2

CAS Number: 14976-57-9 EC Number: 239-055-2

IUPAC Name: (2R)-2-[2-[(1R)-1-(4-Chlorophenylethoxy]ethyl]-1-methyl-2-pyrrolidine fumarate

Description:

H₁-receptor antagonist. Clinically-used antihistamine.

Physical and Chemical Properties:

Batch Molecular Formula: C₂₁H₂₆CINO.C₄H₄O₄

Batch Molecular Weight: 459.97 Physical Appearance: White solid

Minimum Purity: >98%

Batch Molecular Structure:

Storage: Store at RT

Solubility & Usage Info:

DMSO to 50 mM ethanol to 10 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C

water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions

should be made up and used on the same day.

References:

Hartmann et al (1981) Modulation of histamine-induced bronchoconstriction with inhaled, oral, and intravenous clemastine in normal and asthmatic subjects. Thorax 36 737. PMID: 7330790.

Dorsch et al (1982) Histamine1-histamine2 antagonism: effect of combined clemastine and cimetidine pretreatment on allergen and histamine-induced reactions of the guinea pig lung in vivo and in vitro. Agents Actions 12 113. PMID: 6177209.

Nair and Schwartz (1983) Effect of histamine and histamine antagonists on natural and antibody-dependent cellular cytotoxicity of human lymphocytes in vitro. Cell.Immunol. 81 45. PMID: 6225527.

