



Certificate of Analysis

www.tocris.com

Product Name: AICAR Catalog No.: 2840 Batch No.: 6

CAS Number: 2627-69-2 EC Number: 220-097-5

IUPAC Name: N^1 -(β -D-Ribofuranosyl)-5-aminoimidazole-4-carboxamide

1. PHYSICAL AND CHEMICAL PROPERTIES

Storage: Store at -20°C

Batch Molecular Structure:

2. ANALYTICAL DATA

HPLC: Shows 100% purity

¹H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 41.86 5.46 21.69 Found 41.8 5.4 21.57

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use





Product Information

Print Date: Oct 22nd 2013

www.tocris.com

Product Name: AICAR Catalog No.: 2840 Batch No.: 6

CAS Number: 2627-69-2 EC Number: 220-097-5

IUPAC Name: N^1 -(β -D-Ribofuranosyl)-5-aminoimidazole-4-carboxamide

Description:

Cell-permeable, allosteric activator of AMP-activated protein kinase (AMPK). Augments proliferation, differentiation and mineralization of osteoblastic MC3T3-EI cells and attenuates psychosine-induced expression of proinflammatory cytokines and iNOS in astrocytes. Promotes osteogenic differentiation of hAMSCs and BM-MSCs in vitro.

Physical and Chemical Properties:

Batch Molecular Formula: $C_9H_{14}N_4O_5$ Batch Molecular Weight: 258.23 Physical Appearance: White solid

Minimum Purity: >99%

Batch Molecular Structure:

Storage: Store at -20°C

Solubility & Usage Info:

water to 75 mM DMSO to 75 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Giri *et al* (2004) 5-Aminoimidazole-4-carboxamide-1-β-4-ribofuranoside inhibits proinflammatory response in glial cells: a possible role of AMP-activated protein kinase. J.Neurosci. **24** 479. PMID: 14724246.

Kanazawa et al (2007) Adiponectin and AMP kinase activator stimulate proliferation, differentiation and mineralization of osteoblastic MC3T3-EI cells. BMC Cell Biol. 8 51. PMID: 18047638.

Giri et al (2008) The role of AMPK in psychosine mediated effects on oligodendrocytes and astrocytes: Implication for Krabbe disease. J.Neurochem. **105** 1820. PMID: 18248608.

Wu *et al* (2011) AICAR, a small chemical molecule, primes osteogenic differentiation of adult mesenchymal stem cells. Int.J.Artif.Organs. *34* 1128. PMID: 22198598.