

Batch Molecular Formula:

Batch Molecular Weight: Physical Appearance:

Batch Molecular Structure:

Certificate of Analysis

3-[[[2-Methoxy-4-(phenylamino)phenyl]amino]sulfonyl]-2-thiophenecarboxylic acid methyl ester


www.tocris.com

Product Name: GSK 0660 GSK 0660

CAS Number: **IUPAC Name:** 1014691-61-2

Catalog No.: 3433 Batch No.: 1

1. PHYSICAL AND CHEMICAL PROPERTIES C₁₉H₁₈N₂O₅S₂ 418.49 green solid DMSO to 100 mM ethanol to 10 mM



2. ANALYTICAL DATA

Solubility:

Storage:

TLC: HPLC: ¹H NMR: Mass Spectrum: **Microanalysis:**

R_f = 0.7 (Ethyl acetate:Petroleum ether [1:1]) Shows >99.9% purity Consistent with structure Consistent with structure Carbon Hydrogen Nitrogen Theoretical 54.53 4.34 6.69 Found 54.37 4.41 6.65

OMe

ÇO₂Me

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use





Batch No.: 1

www.tocris.com

Product Name: GSK 0660 GSK 0660

CAS Number: 1014691-61-2

3-[[[2-Methoxy-4-(phenylamino)phenyl]amino]sulfonyl]-2-thiophenecarboxylic acid methyl ester

Description:

IUPAC Name:

Selective PPAR δ antagonist (IC₅₀ values are 0.155, > 10 and ≥ 10 µM at PPAR δ , PPAR α and PPAR γ respectively). Exhibits inverse agonist effects when administered by itself.

Physical and Chemical Properties:


Batch Molecular Formula: C₁₉H₁₈N₂O₅S₂

Batch Molecular Weight: 418.49

Physical Appearance: green solid

Minimum Purity: >99%

Batch Molecular Structure:



Storage: Store at +4°C

Solubility & Useage Info:

DMSO to 100 mM ethanol to 10 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a $45-60^{\circ}$ C water bath).

Catalog No.: 3433

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Shearer *et al* (2008) Identification and characterization of a selective peroxisome proliferator-activated receptor β/δ (NR1C2) antagonist. Mol.Endocrinol. **22** 523. PMID: 17975020.

Wang (2008) PPAR-ō in vascular pathophysiology. PPAR Res. 164163. PMID: 19132133.

Bishop-Bailey and Swales (2008) The role of PPARs in the endothelium: Implications for cancer therapy. PPAR Res. 904851.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

