Batch No.: 1



Certificate of Analysis

www.tocris.com

Catalog No.: 3528

Product Name: SCIO 469 hydrochloride

CAS Number: 309913-83-5

IUPAC Name: 6-Chloro-5-[[(2R,5S)-4-[(4-fluorophenyl)methyl]-2,5-dimethyl-1-piperazinyl]carbonyl]-N,N,1-trimethyl- α -oxo-1H-

Indole-3-acetamide hydrochloride

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{27}H_{30}CIFN_4O_3.HCI.^{3}_4H_2O_3$

Batch Molecular Weight: 562.97

Physical Appearance: White solid

Solubility: water to 10 mM

DMSO to 100 mM

Storage: Desiccate at +4°C

Batch Molecular Structure:

2. ANALYTICAL DATA

TLC: $R_f = 0.3$ (Ethanol:Acetic acid)

HPLC: Shows 99.7% purity
Chiral HPLC: Shows 99.3% purity

1H NMR: Consistent with structure
Mass Spectrum: Consistent with structure

Optical Rotation: $[\alpha]_D = +22.9$ (Concentration = 1.04, Solvent = Methanol)

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 57.6 5.82 9.95 Found 57.83 5.77 9.74

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use





Product Information

Print Date: May 18th 2012

Batch No.: 1

www.tocris.com

Product Name: SCIO 469 hydrochloride

CAS Number: 309913-83-5

IUPAC Name: 6-Chloro-5-[[(2R,5S)-4-[(4-fluorophenyl)methyl]-2,5-dimethyl-1-piperazinyl]carbonyl]-N,N,1-trimethyl- α -oxo-1H-

Indole-3-acetamide hydrochloride

Description:

Selective, ATP-competitive p38 inhibitor ($IC_{50} = 9$ nM for p38 α in vitro). Displays approximately 10-fold selectivity for p38 α over p38 β and 2000-fold selectivity for p38 α over 20 other kinases. Reduces p38 α phosphorylation in multiple myeloma cells in vitro and in vivo; activity results in decreased tumor burden and angiogenesis in murine models of multiple myeloma. Also enhances bortezomib-induced cytotoxicity against multiple myeloma cells.

Physical and Chemical Properties:

Batch Molecular Formula: C₂₇H₃₀CIFN₄O₃.HCl.³/₄H₂O

Batch Molecular Weight: 562.97 Physical Appearance: White solid

Minimum Purity: >98%

Batch Molecular Structure:

Storage: Desiccate at +4°C

Solubility & Usage Info:

water to 10 mM DMSO to 100 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Catalog No.: 3528

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Hideshima *et al* (2004) p38 MAPK inhibition enhances PS-341 (bortezomib)-induced cytotoxicity against multiple myeloma cells. Oncogene. **23** 8766. PMID: 15480425.

Giafis et al (2006) Role of the p38 mitogen-activated protein kinase pathway in the generation of arsenic trioxide-dependent cellular responses. Cancer Res. 66 6763. PMID: 16818652.

Vanderkerken *et al* (2007) Inhibition of p38α mitogen-activated protein kinase prevents the development of osteolytic bone disease, reduces tumor burden, and increases survival in murine models of multiple myeloma. Cancer Res. *67* 4572. PMID: 17495322.

RD