



Certificate of Analysis

www.tocris.com

Product Name: TCS 5861528 Catalog No.: 3938 Batch No.: 1

CAS Number: 332117-28-9

IUPAC Name: 2-(1,3-Dimethyl-2,6-dioxo-1,2,3,6-tetrahydro-7*H*-purin-7-yl)-*N*-[4-(1-methylpropyl)phenyl]acetamide

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{19}H_{23}N_5O_3$ Batch Molecular Weight: 369.42

Physical Appearance: white solid

Solubility: DMSO to 100 mM ethanol to 10 mM

Storage: Store at RT

Batch Molecular Structure:

2. ANALYTICAL DATA

HPLC: Shows 98.7% purity

1H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 61.77 6.28 18.96 Found 61.52 6.13 18.9



Product Information

Print Date: Jan 17th 2012

www.tocris.com

Product Name: TCS 5861528 Catalog No.: 3938 Batch No.: 1

CAS Number: 332117-28-9

IUPAC Name: 2-(1,3-Dimethyl-2,6-dioxo-1,2,3,6-tetrahydro-7*H*-purin-7-yl)-*N*-[4-(1-methylpropyl)phenyl]acetamide

Description:

TRPA1 channel blocker that antagonizes AITC- and 4-HNE-evoked calcium influx (IC_{50} values are 14.3 and 18.7 μ M respectively). Attenuates diabetic hypersensitivity in in vivo rat model.

Physical and Chemical Properties:

Batch Molecular Formula: $C_{19}H_{23}N_5O_3$

Batch Molecular Weight: 369.42 Physical Appearance: white solid

Minimum Purity: >98%

Batch Molecular Structure:

Storage: Store at RT

Solubility & Useage Info:

DMSO to 100 mM ethanol to 10 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Wei *et al* (2009) Attenuation of mechanical hypersensitivity by an antagonist of the TRPA1 ion channel in diabetic animals. Anesthesiology *111* 147. PMID: 19512877.

Wei et al (2010) Roles of cutaneous versus spinal TRPA1 channels in mechanical hypersensitivity in the diabetic or mustard oil-treated non-diabetic rat. Neuropharmacology. 58 578. PMID: 20004676.

Wei et al (2010) Spinal TRPA1 ion channels contribute to the cutaneous neurogenic inflammation in the rat. Neurosci.Letts. 479

