

Certificate of Analysis

www.tocris.com

EC Number: 200-037-4

Product Name: 6-Mercaptopurine Catalog No.: 4103 Batch No.: 1

CAS Number: 50-44-2

IUPAC Name: Purine-6(1H)-thione

1. PHYSICAL AND CHEMICAL PROPERTIES

 $C_5H_4N_4S.H_2O$ **Batch Molecular Formula:**

170.2 **Batch Molecular Weight:**

Physical Appearance: Yellow solid

Solubility: DMSO to 100 mM

1eq. NaOH to 100 mM ethanol to 10 mM

Store at RT Storage:

Batch Molecular Structure:

2. ANALYTICAL DATA

HPLC: Shows 99.6% purity ¹H NMR: Consistent with structure **Mass Spectrum:** Consistent with structure

Microanalysis:

Theoretical 35.29 3.55 32.92 Found 35.5 3.58 32.85

Carbon Hydrogen Nitrogen



Product Information

Print Date: Dec 14th 2011

www.tocris.com

Product Name: 6-Mercaptopurine Catalog No.: 4103 Batch No.: 1

CAS Number: 50-44-2 EC Number: 200-037-4

IUPAC Name: Purine-6(1*H*)-thione

Description:

Inhibitor of de novo purine synthesis through interference with DNA and RNA synthesis. Immunosuppressive and antileukemic drug; reduces the anticoagulation elicited by warfarin. Active metabolite of azathioprine (Cat. No. 4099).

Physical and Chemical Properties:

Batch Molecular Formula: $C_5H_4N_4S.H_2O$

Batch Molecular Weight: 170.2 Physical Appearance: Yellow solid

Minimum Purity: >99%

Batch Molecular Structure:

Storage: Store at RT

Solubility & Useage Info:

DMSO to 100 mM 1eq. NaOH to 100 mM ethanol to 10 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Sahasranaman et al (2008) Clinical pharmacology and pharmacogenetics of thiopurines. Eur.J.Clin.Pharmacol. 64 753. PMID: 18506437.

